

# The log-behavior of $\sqrt[n]{p(n)}$ and $\sqrt[n]{p(n)/n}$

William Y.C. Chen<sup>1</sup> and Ken Y. Zheng<sup>2</sup>

<sup>1</sup>Center for Applied Mathematics  
Tianjin University  
Tianjin 300072, P. R. China

<sup>2</sup>Center for Combinatorics, LPMC  
Nankai University  
Tianjin 300071, P. R. China

Email: <sup>1</sup>chenyc@tju.edu.cn, <sup>2</sup>kenzheng@aliyun.com

## Abstract

Let  $p(n)$  denote the partition function and let  $\Delta$  be the difference operator respect to  $n$ . In this paper, we obtain a lower bound for  $\Delta^2 \log \sqrt[n-1]{p(n-1)/(n-1)}$ , leading to a proof of the conjecture of Sun on the log-convexity of  $\{\sqrt[n]{p(n)/n}\}_{n \geq 60}$ . Using the same argument, it can be shown that for any real number  $\alpha$ , there exists an integer  $n(\alpha)$  such that the sequence  $\{\sqrt[n]{p(n)/n^\alpha}\}_{n \geq n(\alpha)}$  is log-convex. Moreover, we show that  $\lim_{n \rightarrow +\infty} n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)} = 3\pi/\sqrt{24}$ . Finally, by finding an upper bound of  $\Delta^2 \log \sqrt[n-1]{p(n-1)}$ , we establish an inequality on the ratio  $\frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}$ .

**Keywords:** Partition function, Log-convex sequence, Hardy-Ramanujan-Rademacher formula, Lehmer's error bound

**AMS Subject Classifications:** 05A20

## 1 Introduction

The objective of this paper is to study the log-behavior of the sequences  $\sqrt[n]{p(n)}$  and  $\sqrt[n]{p(n)/n}$ , where  $p(n)$  denotes the number of partitions of a nonnegative integer  $n$ . A positive sequence  $\{a_n\}_{n \geq 0}$  is log-convex if it satisfies that for  $n \geq 1$ ,

$$a_n^2 - a_{n-1}a_{n+1} \leq 0,$$

and it is called log-concave if for  $n \geq 1$ ,

$$a_n^2 - a_{n-1}a_{n+1} \geq 0.$$

Let  $r(n) = \sqrt[n]{p(n)/n}$  and let  $\Delta$  be the difference operator respect to  $n$ . Sun [11] conjectured that the sequence  $\{r(n)\}_{n \geq 60}$  is log-convex. Desalvo and Pak [5] noticed that the log-convexity of  $\{r(n)\}_{n \geq 60}$  can be derived from an estimate for  $\Delta^2 \log r(n-1)$ , see [5, Final Remark 7.7]. They also remarked that their approach to bounding  $-\Delta^2 \log p(n-1)$  does not seem to apply to  $\Delta^2 \log r(n-1)$ . In this paper, we obtain a lower bound for  $\Delta^2 \log r(n-1)$ , leading to a proof of the log-convexity of  $\{r(n)\}_{n \geq 60}$ .

**Theorem 1.1** *The sequence  $\{r(n)\}_{n \geq 60}$  is log-convex.*

The log-convexity of  $\{r(n)\}_{n \geq 60}$  implies the log-convexity of  $\{\sqrt[n]{p(n)}\}_{n \geq 26}$ , because the sequence  $\{\sqrt[n]{n}\}_{n \geq 4}$  is log-convex [11]. It is known that  $\lim_{n \rightarrow +\infty} \sqrt[n]{p(n)} = 1$ . For a combinatorial proof of this fact, see Andrews [1]. Sun [11] proposed the conjecture that  $\{\sqrt[n]{p(n)}\}_{n \geq 6}$  is strictly decreasing, which has been proved by Wang and Zhu [12]. The log-convexity of  $\{\sqrt[n]{p(n)}\}_{n \geq 26}$  was also conjectured by Sun [11]. It is easy to see that the log-convexity of  $\{\sqrt[n]{p(n)}\}_{n \geq 26}$  implies the decreasing property.

It should be noted that there is an alternative way to prove the log-convexity of  $\{\sqrt[n]{p(n)}\}_{n \geq 26}$ . Chen, Guo and Wang [3] introduced the notion of a ratio log-convex sequence and showed that the ratio log-convexity implies the log-convexity under a certain initial condition. A sequence  $\{a_n\}_{n \geq k}$  is called ratio log-convex if  $\{a_{n+1}/a_n\}_{n \geq k}$  is log-convex, or, equivalently, for  $n \geq k+1$ ,

$$\log a_{n+2} - 3 \log a_{n+1} + 3 \log a_n - \log a_{n-1} \geq 0.$$

Chen et al. [4] showed that that for any  $r \geq 1$ , one can determine a number  $n(r)$  such that for  $n > n(r)$ ,  $(-1)^{r-1} \Delta^r \log p(n)$  is positive. For  $r = 3$ , it can be shown that for  $n \geq 116$ ,

$$\Delta^3 \log p(n-1) > 0.$$

Since

$$\Delta^3 \log p(n-1) = \log p(n+2) - 3 \log p(n+1) + 3 \log p(n) - \log p(n-1),$$

we see  $\{p(n)\}_{n \geq 115}$  is ratio log-convex. So we are led to the following assertion.

**Theorem 1.2** *The sequence  $\{\sqrt[n]{p(n)}\}_{n \geq 26}$  is log-convex.*

Moreover, as pointed out by the referee, we may consider the log-behavior of  $\sqrt[n]{p(n)/n^\alpha}$  for any real number  $\alpha$ . To this end, we obtain the following generalization of Theorems 1.1 and 1.2.

**Theorem 1.3** *Let  $\alpha$  be a real number. There exists a positive integer  $n(\alpha)$  such that the sequence  $\{\sqrt[n]{p(n)/n^\alpha}\}_{n \geq n(\alpha)}$  is log-convex.*

We also establish the following inequality on the ratio  $\frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}$ .

**Theorem 1.4** *For  $n \geq 2$ , we have*

$$\frac{\sqrt[n]{p(n)}}{\sqrt[n+1]{p(n+1)}} \left( 1 + \frac{3\pi}{\sqrt{24}n^{5/2}} \right) > \frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}. \quad (1.1)$$

Desalvo and Pak [5] have shown that the limit of  $-n^{\frac{3}{2}}\Delta^2 \log p(n)$  is  $\pi/\sqrt{24}$ . By bounding  $\Delta^2 \log \sqrt[n]{p(n)}$ , we derive the following limit of  $n^{\frac{5}{2}}\Delta^2 \log \sqrt[n]{p(n)}$ :

$$\lim_{n \rightarrow +\infty} n^{\frac{5}{2}}\Delta^2 \log \sqrt[n]{p(n)} = 3\pi/\sqrt{24}. \quad (1.2)$$

From the above relation (1.2), it can be seen that the coefficient  $\frac{3\pi}{\sqrt{24}}$  in (1.1) is the best possible.

## 2 The Log-convexity of $r(n)$

In this section, we obtain a lower bound of  $\Delta^2 \log r(n-1)$  and prove the log-convexity of  $\{r(n)\}_{n \geq 60}$ . First, we follow the approach of Desalvo and Pak to give an expression of  $\Delta^2 \log r(n-1)$  as a sum of  $\Delta^2 \tilde{B}(n-1)$  and  $\Delta^2 \tilde{E}(n-1)$ , where  $\Delta^2 \tilde{B}(n-1)$  makes a major contribution to  $\Delta^2 \log r(n-1)$  with  $\Delta^2 \tilde{E}(n-1)$  being the error term, that is,  $\Delta^2 \tilde{B}(n-1)$  converges to  $\Delta^2 \log r(n-1)$ . The expressions for  $B(n)$  and  $E(n)$  will be given later. In this setting, we derive a lower bound of  $\Delta^2 \tilde{B}(n-1)$ . By Lehmer's error bound, we give an upper bound for  $|\Delta^2 \tilde{E}(n-1)|$ . Combining the lower bound for  $\Delta^2 \tilde{B}(n-1)$  and the upper bound for  $\Delta^2 \tilde{E}(n-1)$ , we are led to a lower bound for  $\Delta^2 \log r(n-1)$ . By proving the positivity of this lower bound for  $\Delta^2 \log r(n-1)$ , we reach the log-convexity of  $\{r(n)\}_{n \geq 60}$ .

The strict log-convexity of  $\{r(n)\}_{n \geq 60}$  can be restated as the following relation for  $n \geq 61$

$$\log r(n+1) + \log r(n-1) - 2\log r(n) > 0,$$

that is, for  $n \geq 61$ ,

$$\Delta^2 \log r(n-1) > 0.$$

For  $n \geq 1$  and any positive integer  $N$ , the Hardy-Ramanujan-Rademacher formula (see [2, 6, 7, 10]) reads

$$p(n) = \frac{d}{\mu^2} \sum_{k=1}^N A_k^*(n) \left[ \left( 1 - \frac{k}{\mu} \right) e^{\frac{\mu}{k}} + \left( 1 + \frac{k}{\mu} \right) e^{-\frac{\mu}{k}} \right] + R_2(n, N), \quad (2.1)$$

where  $d = \frac{\pi^2}{6\sqrt{3}}$ ,  $\mu(n) = \frac{\pi}{6}\sqrt{24n-1}$ ,  $A_k^*(n) = k^{-\frac{1}{2}}A_k(n)$ ,  $A_k(n)$  is a sum of 24th roots of unity with initial values  $A_1(n) = 1$  and  $A_2(n) = (-1)^n$ ,  $R_2(n, N)$  is the remainder. Lehmer's error bound (see [8, 9]) for  $R_2(n, N)$  is given by

$$|R_2(n, N)| < \frac{\pi^2 N^{-2/3}}{\sqrt{3}} \left[ \left( \frac{N}{\mu} \right)^3 \sinh \frac{\mu}{N} + \frac{1}{6} - \left( \frac{N}{\mu} \right)^2 \right]. \quad (2.2)$$

Let us give an outline of Desalvo and Pak's approach to proving the log-concavity of  $\{p(n)\}_{n>25}$ . Setting  $N = 2$  in (2.1), they expressed  $p(n)$  as

$$p(n) = T(n) + R(n), \quad (2.3)$$

where

$$T(n) = \frac{d}{\mu(n)^2} \left[ \left( 1 - \frac{1}{\mu(n)} \right) e^{\mu(n)} + \frac{(-1)^n}{\sqrt{2}} e^{\frac{\mu(n)}{2}} \right], \quad (2.4)$$

$$R(n) = \frac{d}{\mu(n)^2} \left[ \left( 1 + \frac{1}{\mu(n)} \right) e^{-\mu(n)} - \frac{(-1)^n}{\sqrt{2}} \frac{2}{\mu(n)} + \frac{(-1)^n}{\sqrt{2}} \left( 1 + \frac{2}{\mu(n)} \right) e^{-\frac{\mu(n)}{2}} \right] + R_2(n, 2). \quad (2.5)$$

They have shown that

$$\left| \Delta^2 \log p(n-1) - \Delta^2 \log T(n-1) \right| = \left| \Delta^2 \log \left( 1 + \frac{R(n-1)}{T(n-1)} \right) \right| < e^{-\frac{\pi\sqrt{2n}}{10\sqrt{3}}} \quad (2.6)$$

and

$$\left| \Delta^2 \log T(n-1) - \Delta^2 \log \frac{d}{\mu(n-1)^2} \left( 1 - \frac{1}{\mu(n-1)} \right) e^{\mu(n-1)} \right| < e^{-\frac{\pi\sqrt{2n}}{10\sqrt{3}}}. \quad (2.7)$$

It follows that  $\Delta^2 \log \frac{d}{\mu(n-1)^2} \left( 1 - \frac{1}{\mu(n-1)} \right) e^{\mu(n-1)}$  converges to  $\Delta^2 \log p(n-1)$ . Finally, they use  $-\Delta^2 \log \frac{d}{\mu(n-1)^2} \left( 1 - \frac{1}{\mu(n-1)} \right) e^{\mu(n-1)}$  to estimate  $-\Delta^2 \log p(n-1)$ , leading to the log-concavity of  $\{p(n)\}_{n>25}$ .

We shall use an alternative decomposition of  $p(n)$ . Setting  $N = 2$  in (2.1), we can express  $p(n)$  as

$$p(n) = \tilde{T}(n) + \tilde{R}(n), \quad (2.8)$$

where

$$\tilde{T}(n) = \frac{d}{\mu(n)^2} \left( 1 - \frac{1}{\mu(n)} \right) e^{\mu(n)}, \quad (2.9)$$

$$\begin{aligned} \tilde{R}(n) = & \frac{d}{\mu(n)^2} \left[ \left( 1 + \frac{1}{\mu(n)} \right) e^{-\mu(n)} + \frac{(-1)^n}{\sqrt{2}} \left( 1 - \frac{2}{\mu(n)} \right) e^{\frac{\mu(n)}{2}} \right. \\ & \left. + \frac{(-1)^n}{\sqrt{2}} \left( 1 + \frac{2}{\mu(n)} \right) e^{-\frac{\mu(n)}{2}} \right] + R_2(n, 2). \end{aligned} \quad (2.10)$$

Based on the decomposition (2.8) for  $p(n)$ , one can express  $\Delta^2 \log r(n-1)$  as follows:

$$\Delta^2 \log r(n-1) = \Delta^2 \tilde{B}(n-1) + \Delta^2 \tilde{E}(n-1), \quad (2.11)$$

where

$$\tilde{B}(n) = \frac{1}{n} \log \tilde{T}(n) - \frac{1}{n} \log n, \quad (2.12)$$

$$\tilde{y}_n = \tilde{R}(n)/\tilde{T}(n), \quad (2.13)$$

$$\tilde{E}(n) = \frac{1}{n} \log(1 + \tilde{y}_n). \quad (2.14)$$

The following lemma will be used to derive a lower bound and an upper bound of  $\Delta^2 \tilde{B}(n-1)$ .

**Lemma 2.1** *Suppose  $f(x)$  has a continuous second derivative for  $x \in [n-1, n+1]$ . Then there exists  $c \in (n-1, n+1)$  such that*

$$\Delta^2 f(n-1) = f(n+1) + f(n-1) - 2f(n) = f''(c). \quad (2.15)$$

*If  $f(x)$  has an increasing second derivative, then*

$$f''(n-1) < \Delta^2 f(n-1) < f''(n+1). \quad (2.16)$$

*Conversely, if  $f(x)$  has a decreasing second derivative, then*

$$f''(n+1) < \Delta^2 f(n-1) < f''(n-1). \quad (2.17)$$

*Proof.* Set  $\varphi(x) = f(x+1) - f(x)$ . By the mean value theorem, there exists a number  $\xi \in (n-1, n)$  such that

$$f(n+1) + f(n-1) - 2f(n) = \varphi(n) - \varphi(n-1) = \varphi'(\xi).$$

Again, applying the mean value theorem to  $\varphi'(\xi)$ , there exists a number  $\theta \in (0, 1)$  such that

$$\varphi'(\xi) = f'(\xi+1) - f'(\xi) = f''(\xi+\theta).$$

Let  $c = \xi + \theta$ . Then we get (2.15), which yields (2.16) and (2.17). ■

In order to find a lower bound for  $\Delta^2 \log r(n-1)$  and obtain the limit of  $n^{\frac{5}{2}} \Delta^2 \log \sqrt[n]{p(n)}$ , we need the following lower and upper bounds for  $\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1)$ .

**Lemma 2.2** *Let*

$$B_1(n) = \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{4\log(\mu(n-1))}{(n-1)^3}, \quad (2.18)$$

$$B_2(n) = \frac{72\pi}{(n-1)(24n-25)^{3/2}} - \frac{4\log(\mu(n+1))}{(n+1)^3} + \frac{5}{(n-1)^3}. \quad (2.19)$$

For  $n \geq 40$ , we have

$$B_1(n) < \Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) < B_2(n). \quad (2.20)$$

*Proof.* By the definition (2.9), we may write

$$\frac{\log \tilde{T}(n)}{n} = \sum_{i=1}^4 f_i,$$

where

$$\begin{aligned} f_1(n) &= \frac{\mu(n)}{n}, \\ f_2(n) &= -\frac{3\log \mu(n)}{n}, \\ f_3(n) &= \frac{\log(\mu(n)-1)}{n}, \\ f_4(n) &= \frac{\log d}{n}. \end{aligned}$$

Thus

$$\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) = \sum_{i=1}^4 \Delta^2 f_i(n-1). \quad (2.21)$$

Since

$$f_1'''(n) = \frac{\pi}{n(24n-1)^{3/2}} \left( -\frac{216}{n} + \frac{864}{24n-1} + \frac{36}{n^2} - \frac{1}{n^3} \right),$$

we see that for  $n \geq 1$ ,  $f_1'''(n) < 0$ . Similarly, it can be checked that for  $n \geq 4$ ,  $f_2'''(n) > 0$ ,  $f_3'''(n) < 0$ , and  $f_4'''(n) > 0$ . Consequently, for  $n \geq 4$ ,  $f_1''(n)$  and  $f_3''(n)$  are decreasing, whereas  $f_2''(n)$  and  $f_4''(n)$  are increasing. Using Lemma 2.1, for each  $i$ , we can get a lower bound and an upper bound for  $\Delta^2 f_i(n-1)$  in terms of  $f_i''(n-1)$  and  $f_i''(n+1)$ . For example,

$$f_1''(n+1) < \Delta^2 f_1(n-1) < f_1''(n-1).$$

So, by (2.21) we find that

$$\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) > f_1''(n+1) + f_2''(n-1) + f_3''(n+1) + f_4''(n-1), \quad (2.22)$$

and

$$\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) < f_1''(n-1) + f_2''(n+1) + f_3''(n-1) + f_4''(n+1), \quad (2.23)$$

where

$$f_1''(n) = \frac{72\pi}{n(24n-1)^{3/2}} - \frac{12\pi}{n^2(24n-1)^{3/2}} + \frac{\pi}{3n^3(24n-1)^{3/2}}, \quad (2.24)$$

$$f_2''(n) = -\frac{6 \log \mu(n)}{n^3} + \frac{72}{(24n-1)n^2} + \frac{864}{n(24n-1)^2}, \quad (2.25)$$

$$f_3''(n) = -\frac{4\pi^2}{(\mu(n)-1)^2(24n-1)n} + \frac{2 \log(\mu(n)-1)}{n^3} \\ - \frac{4\pi}{(\mu(n)-1)\sqrt{24n-1}n^2} - \frac{24\pi}{(\mu(n)-1)(24n-1)^{3/2}n}, \quad (2.26)$$

$$f_4''(n) = \frac{2 \log d}{n^3}. \quad (2.27)$$

According to (2.24), one can check that for  $n \geq 2$ ,

$$f_1''(n+1) > \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{12\pi}{(n+1)^2(24n+23)^{3/2}}. \quad (2.28)$$

An easy computation shows that for  $n \geq 3$ ,

$$\mu(n) - 1 > \frac{2}{3}\mu(n-2). \quad (2.29)$$

Substituting (2.29) into (2.26) yields that

$$f_3''(n+1) > \frac{2 \log(\mu(n+1)-1)}{(n+1)^3} - \frac{540}{(24n-25)^2(n-1)} - \frac{36}{(24n-25)(n-1)^2}. \quad (2.30)$$

Using (2.25) and (2.30), we find that

$$f_2''(n-1) + f_3''(n+1) \\ > \frac{2 \log(\mu(n+1)-1)}{(n+1)^3} - \frac{6 \log(\mu(n-1))}{(n-1)^3} \\ + \frac{324}{(n-1)(24n-25)^2} + \frac{36}{(n-1)^2(24n-25)} \quad (2.31)$$

Apparently, for  $n \geq 2$ ,

$$\frac{2}{(n+1)^3} - \frac{2}{(n-1)^3} > -\frac{12}{(n-1)^4},$$

so that

$$\begin{aligned}
& \frac{2 \log(\mu(n+1) - 1)}{(n+1)^3} - \frac{6 \log(\mu(n-1))}{(n-1)^3} \\
& > \frac{2 \log(\mu(n+1) - 1)}{(n+1)^3} - \frac{2 \log(\mu(n+1) - 1)}{(n-1)^3} - \frac{4 \log(\mu(n-1))}{(n-1)^3} \\
& > -\frac{12 \log(\mu(n+1) - 1)}{(n-1)^4} - \frac{4 \log(\mu(n-1))}{(n-1)^3}.
\end{aligned} \tag{2.32}$$

Since, for  $n \geq 2$ ,

$$\frac{324}{(n-1)(24n-25)^2} + \frac{36}{(n-1)^2(24n-25)} > \frac{2}{(n-1)^3}, \tag{2.33}$$

utilizing (2.31) and (2.32) yields, for  $n \geq 3$ ,

$$f_2''(n-1) + f_3''(n+1) > -\frac{4 \log(\mu(n-1))}{(n-1)^3} + \frac{2}{(n-1)^3} - \frac{12 \log(\mu(n+1) - 1)}{(n-1)^4}. \tag{2.34}$$

Using (2.27), (2.28) and (2.34), we deduce that

$$\begin{aligned}
& f_1''(n+1) + f_2''(n-1) + f_3''(n+1) + f_4''(n-1) - B_1(n) \\
& > \frac{2(1 + \log d)}{(n-1)^3} - \frac{12\pi}{(n+1)^2(24n+23)^{3/2}} - \frac{12 \log(\mu(n+1) - 1)}{(n-1)^4}.
\end{aligned} \tag{2.35}$$

Let  $C(n)$  be the right hand side of (2.35). By (2.22), to prove  $B_1(n) < \Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1)$ , it is enough to show that  $C(n) > 0$  when  $n \geq 40$ . Since  $\log x < x$  for  $x > 0$ , for  $n \geq 3$

$$\mu(n+1) - 1 < \frac{\pi}{4} \sqrt{24n-24}, \tag{2.36}$$

we get

$$-\frac{12 \log(\mu(n+1) - 1)}{(n-1)^4} > -\frac{12(\mu(n+1) - 1)}{(n-1)^4} > -\frac{3\sqrt{24}\pi}{(n-1)^{7/2}}. \tag{2.37}$$

Note that for  $n \geq 2$ ,

$$-\frac{12\pi}{(n+1)^2(24n+23)^{3/2}} > -\frac{\sqrt{24}\pi}{48(n-1)^{7/2}}. \tag{2.38}$$

Combining (2.37) and (2.38), we see that for  $n \geq 2$ ,

$$C(n) > \frac{2(1 + \log d)}{(n-1)^3} - \frac{(3 + 1/48)\sqrt{24}\pi}{(n-1)^{7/2}}. \tag{2.39}$$

It is straightforward to show that the right hand side of (2.39) is positive if  $n \geq 490$ . For  $40 \leq n \leq 489$ , it is routine to check that  $C(n) > 0$ , and so  $C(n) > 0$  for  $n \geq 40$ . It follows from (2.35) that for  $n \geq 40$ ,

$$\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) > B_1(n).$$



To derive the upper bound for  $\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1)$ , we obtain the following upper bounds which can be verified directly. The proofs are omitted. For  $n \geq 2$ ,

$$\begin{aligned}
f_1''(n-1) &< \frac{72\pi}{(n-1)[24n-25]^{3/2}}, \\
f_2''(n+1) &< -\frac{6 \log \mu(n+1)}{(n+1)^3} + \frac{9}{2(n-1)^3}, \\
f_3''(n-1) &< -\frac{4\pi^2}{(\mu(n-1))^2(24n-25)(n-1)} + \frac{2 \log(\mu(n-1))}{(n-1)^3} \\
&\quad - \frac{4\pi}{\mu(n-1)\sqrt{24n-25}(n-1)^2} - \frac{24\pi}{\mu(n-1)(24n-25)^{3/2}(n-1)}, \\
f_2''(n+1) + f_3''(n-1) &< \frac{3}{(n-1)^3} + \frac{12 \log(\mu(n+1))}{(n-1)^4} - \frac{4 \log(\mu(n+1))}{(n+1)^3}, \\
f_4''(n+1) &< 0.
\end{aligned}$$

Combining the above upper bounds, we conclude that for  $n \geq 40$ ,

$$f_1''(n-1) + f_2''(n+1) + f_3''(n-1) + f_4''(n+1) < B_2(n).$$

This completes the proof. ■

The following lemma gives an upper bound for  $|\Delta^2 \tilde{E}(n-1)|$ .

**Lemma 2.3** For  $n \geq 40$ ,

$$|\Delta^2 \tilde{E}(n-1)| < \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \quad (2.40)$$

*Proof.* By (2.14), we find that for  $n \geq 2$ ,

$$\Delta^2 \tilde{E}(n-1) = \frac{1}{n-1} \log(1 + \tilde{y}_{n-1}) + \frac{1}{n+1} \log(1 + \tilde{y}_{n+1}) - \frac{2}{n} \log(1 + \tilde{y}_n), \quad (2.41)$$

where

$$\tilde{y}_n = \tilde{R}(n)/\tilde{T}(n).$$

To bound  $|\Delta^2 \tilde{E}(n-1)|$ , it is necessary to bound  $\tilde{y}_n$ . For this purpose, we first consider  $\tilde{R}(n)$ , as defined by (2.10). Since  $d < 1$  and  $\mu(n) > 2$ , for  $n \geq 1$  we have

$$\begin{aligned}
&\frac{d}{\mu(n)^2} \left[ \left(1 + \frac{1}{\mu(n)}\right) e^{-\mu(n)} + \frac{(-1)^n}{\sqrt{2}} \left(1 - \frac{2}{\mu(n)}\right) e^{\frac{\mu(n)}{2}} + \frac{(-1)^n}{\sqrt{2}} \left(1 + \frac{2}{\mu(n)}\right) e^{-\frac{\mu(n)}{2}} \right] \\
&< \frac{1}{\mu(n)^2} \left(1 + e^{\frac{\mu(n)}{2}} + 1\right).
\end{aligned}$$

For  $N = 2$  and  $n \geq 1$ , Lehmer's bound (2.2) reduces to

$$|R_2(n, 2)| < 4 \left( 1 + \frac{4}{\mu(n)^3} e^{\frac{\mu(n)}{2}} \right).$$

By the definition of  $\tilde{R}(n)$ ,

$$|\tilde{R}(n)| < \frac{1}{\mu(n)^2} \left( 1 + e^{\frac{\mu(n)}{2}} + 1 \right) + 4 \left( 1 + \frac{4}{\mu(n)^3} e^{\frac{\mu(n)}{2}} \right) < 5 + \frac{9}{\mu(n)^2} e^{\frac{\mu(n)}{2}}. \quad (2.42)$$

Recalling the definition (2.9) of  $\tilde{T}(n)$ , it follows from (2.42) that for  $n \geq 1$ ,

$$|\tilde{y}_n| < \frac{\mu(n)}{d(\mu(n) - 1)} \left( 5\mu(n)^2 e^{-\frac{2\mu(n)}{3}} + 9e^{-\frac{\mu(n)}{6}} \right) e^{-\frac{\mu(n)}{3}}. \quad (2.43)$$

Observe that for  $n \geq 2$ ,

$$\left( 5\mu(n)^2 e^{-\frac{2\mu(n)}{3}} + 9e^{-\frac{\mu(n)}{6}} \right)' < 0, \quad (2.44)$$

and

$$\left( \frac{d(\mu(n) - 1)}{\mu(n)} \right)' > 0. \quad (2.45)$$

Since

$$5\mu^2(40) e^{-\frac{2\mu(40)}{3}} + 9e^{-\frac{\mu(40)}{6}} < \frac{d(\mu(40) - 1)}{\mu(40)},$$

using (2.44) and (2.45), we deduce that for  $n \geq 40$ ,

$$5\mu^2(n) e^{-\frac{2\mu(n)}{3}} + 9e^{-\frac{\mu(n)}{6}} < \frac{d(\mu(n) - 1)}{\mu(n)}. \quad (2.46)$$

Now, it is clear from (2.43) and (2.46) that for  $n \geq 40$ ,

$$|\tilde{y}_n| < e^{-\frac{\mu(n)}{3}}. \quad (2.47)$$

In view of (2.47), for  $n \geq 40$ ,

$$|\tilde{y}_n| < e^{-\frac{\mu(40)}{3}} < \frac{1}{5}. \quad (2.48)$$

It is known that  $\log(1 + x) < x$  for  $0 < x < 1$  and  $-\log(1 + x) < -x/(1 + x)$  for  $-1 < x < 0$ . Thus, for  $|x| < 1$ ,

$$|\log(1 + x)| \leq \frac{|x|}{1 - |x|}, \quad (2.49)$$

see also [5], and so it follows from (2.48) and (2.49) that for  $n \geq 40$ ,

$$|\log(1 + \tilde{y}_n)| \leq \frac{|\tilde{y}_n|}{1 - |\tilde{y}_n|} \leq \frac{5}{4} |\tilde{y}_n|. \quad (2.50)$$

Because of (2.41), we see that for  $n \geq 2$ ,

$$\left| \Delta^2 \tilde{E}(n-1) \right| \leq \frac{1}{n-1} |\log(1 + \tilde{y}_{n-1})| + \frac{1}{n+1} |\log(1 + \tilde{y}_{n+1})| + \frac{2}{n} |\log(1 + \tilde{y}_n)|. \quad (2.51)$$

Applying (2.50) to (2.51), we obtain that for  $n \geq 40$ ,

$$\left| \Delta^2 \tilde{E}(n-1) \right| \leq \frac{5}{4} \left( \frac{|\tilde{y}_{n-1}|}{n-1} + \frac{|\tilde{y}_{n+1}|}{n+1} + \frac{2|\tilde{y}_n|}{n} \right). \quad (2.52)$$

Plugging (2.47) into (2.52), we infer that for  $n \geq 40$ ,

$$\left| \Delta^2 \tilde{E}(n-1) \right| < \frac{5}{4} \left( \frac{e^{-\frac{\mu(n-1)}{3}}}{n-1} + \frac{e^{-\frac{\mu(n+1)}{3}}}{n+1} + \frac{2e^{-\frac{\mu(n)}{3}}}{n} \right). \quad (2.53)$$

But  $\frac{1}{n} e^{-\frac{\mu(n)}{3}}$  is decreasing for  $n \geq 1$ . It follows from (2.53) that for  $n \geq 40$ ,

$$\left| \Delta^2 \tilde{E}(n-1) \right| < \frac{5}{n-1} e^{-\frac{\mu(n-1)}{3}}.$$

This proves (2.40). ■

With the aid of Lemma 2.2 and 2.3, we are ready to prove the log-convexity of  $\{r(n)\}_{n \geq 60}$ .

*Proof of Theorem 1.1.* To prove the strict log-convexity of  $\{r(n)\}_{n \geq 60}$ , we proceed to show that for  $n \geq 61$ ,

$$\Delta^2 \log r(n-1) > 0.$$

Evidently, for  $n \geq 40$ ,

$$\left( -\frac{\log n}{n} \right)''' > 0.$$

By Lemma 2.1,

$$-\Delta^2 \frac{\log(n-1)}{n-1} > \left( -\frac{\log(n-1)}{n-1} \right)'' ,$$

that is,

$$-\Delta^2 \frac{\log(n-1)}{n-1} > -\frac{2\log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3}. \quad (2.54)$$

It follows from (2.12) that

$$\Delta^2 \tilde{B}(n-1) = \Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) - \Delta^2 \frac{\log(n-1)}{n-1}.$$

Applying Lemma 2.2 and (2.54) to the above relation, we deduce that for  $n \geq 40$ ,

$$\Delta^2 \tilde{B}(n-1) > \tilde{B}_1(n) - \frac{2 \log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3},$$

that is,

$$\Delta^2 \tilde{B}(n-1) > \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2 \log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3}. \quad (2.55)$$

By (2.11) and Lemma 2.3, we find that for  $n \geq 40$ ,

$$\Delta^2 \log r(n-1) > \Delta^2 \tilde{B}(n-1) - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \quad (2.56)$$

It follows from (2.55) and (2.56) that for  $n \geq 40$ ,

$$\begin{aligned} & \Delta^2 \log r(n-1) \\ & > \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2 \log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \end{aligned}$$

Let  $D(n)$  denote the right hand side of the above relation. Clearly, for  $n \geq 5505$ ,

$$\frac{72\pi}{(n+1)(24n+23)^{3/2}} > \frac{3\pi}{\sqrt{24}(n+1)^{5/2}} > \frac{1}{(n-1)^{5/2}}. \quad (2.57)$$

To prove that  $D(n) > 0$  for  $n \geq 5505$ , we wish to show that for  $n \geq 5505$ ,

$$-\frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2 \log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} > -\frac{1}{(n-1)^{5/2}}. \quad (2.58)$$

Using the fact that for  $x > 5504$ ,  $\log x < x^{1/4}$ , we deduce that for  $n \geq 5505$ ,

$$\frac{4 \log[\mu(n-1)]}{(n-1)^3} < \frac{4\sqrt[4]{\mu(n-1)}}{(n-1)^3} < \frac{4\sqrt[4]{\frac{\pi}{4}\sqrt{24n-24}}}{(n-1)^3} < \frac{6}{(n-1)^{23/8}}, \quad (2.59)$$

and

$$\frac{2 \log(n-1)}{(n-1)^3} < \frac{2(n-1)^{1/4}}{(n-1)^3} < \frac{2}{(n-1)^{11/4}}. \quad (2.60)$$

Since  $e^x > x^6/720$  for  $x > 0$ , we see that for  $n \geq 2$ ,

$$\frac{1}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} < \frac{1}{n-1} e^{-\frac{\pi\sqrt{23n}}{18}} < \frac{2094}{n^3(n-1)} < \frac{2094}{(n-1)^4}. \quad (2.61)$$

Combining (2.59), (2.60) and (2.61), we find that for  $n \geq 5505$ ,

$$-\frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2 \log(n-1)}{(n-1)^3} + \frac{3}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}$$

$$\begin{aligned}
&> -\frac{6}{(n-1)^{23/8}} - \frac{2}{(n-1)^{11/4}} + \frac{3}{(n-1)^3} - \frac{10470}{(n-1)^4} \\
&> -\frac{6}{(n-1)^{23/8}} - \frac{2}{(n-1)^{11/4}} \\
&> -\frac{1}{(n-1)^{5/2}}.
\end{aligned}$$

This proves the inequality (2.58). By (2.58) and (2.57), we obtain that  $D(n) > 0$  for  $n \geq 5505$ . Verifying that  $\Delta^2 \log r(n-1) > 0$  for  $61 \leq n \leq 5504$  completes the proof. ■

Clearly, Theorem 1.3 is a generalization as well as a unification of Theorem 1.1 and 1.2. In fact, it can be proved in the same manner as the proof of Theorem 1.1.

*Proof of Theorem 1.3.* Let  $\alpha$  be a real number. When  $\alpha \leq 0$ , it is clear that  $\frac{1}{\sqrt[n]{n^\alpha}}$  is log-convex. It follows from Theorem 1.2 that  $\sqrt[n]{p(n)/n^\alpha}$  is log-convex for  $n \geq 26$ .

We now consider the case  $\alpha > 0$ . A similar argument to the proof of Theorem 1.1 shows that for  $n \geq 40$ ,

$$\begin{aligned}
&\Delta^2 \log \sqrt[n-1]{p(n-1)/(n-1)^\alpha} \\
&= \Delta^2 \frac{1}{n-1} \log T(n) + \Delta^2 \frac{1}{n-1} \log(1 + y_{n-1}) - \alpha \Delta^2 \frac{\log(n-1)}{n-1} \\
&> \frac{72\pi}{(n+1)(24n+23)^{3/2}} - \frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2\alpha \log(n-1)}{(n-1)^3} \\
&\quad + \frac{3\alpha}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \tag{2.62}
\end{aligned}$$

It is easy to check that for  $n \geq \max\left\{\left\lceil\frac{3490}{\alpha}\right\rceil + 2, 5505\right\}$ ,

$$\frac{3\alpha}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} > \frac{3\alpha}{(n-1)^3} - \frac{10470}{(n-1)^4} > 0,$$

and that for  $n \geq \max\{[(2\alpha+3)^4] + 2, 5505\}$ ,

$$-\frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2\alpha \log(n-1)}{(n-1)^3} > -\frac{6}{(n-1)^{23/8}} - \frac{2\alpha}{(n-1)^{11/4}} > -\frac{1}{(n-1)^{5/2}}.$$

Let

$$n(\alpha) = \max\left\{\left\lceil\frac{3490}{\alpha}\right\rceil + 2, [(2\alpha+3)^4] + 2, 5505\right\}.$$

It can be seen that for  $n > n(\alpha)$ ,

$$-\frac{4 \log[\mu(n-1)]}{(n-1)^3} - \frac{2\alpha \log(n-1)}{(n-1)^3} + \frac{3\alpha}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} > -\frac{1}{(n-1)^{5/2}}. \tag{2.63}$$

Combing (2.57) and (2.63), we deduce that the right hand side of (2.62) is positive for  $n > n(\alpha)$ . So we are led to the log-convexity of the sequence  $\{\sqrt[n]{p(n)/n^\alpha}\}_{n \geq n(\alpha)}$ . ■

### 3 An inequality on the ratio $\frac{n^{-1}\sqrt[p(n-1)]{p(n-1)}}{\sqrt[p(n)]{p(n)}}$

In this section, we employ Lemma 2.2 and Lemma 2.3 to find the limit of  $n^{\frac{5}{2}}\Delta^2 \log \sqrt[p(n)]{p(n)}$ . Then we give an upper bound for  $\Delta^2 \log \sqrt[p(n-1)]{p(n-1)}$ . This leads to the inequality (1.1).

**Theorem 3.1** *Let  $\beta = 3\pi/\sqrt{24}$ . We have*

$$\lim_{n \rightarrow +\infty} n^{\frac{5}{2}}\Delta^2 \log \sqrt[p(n)]{p(n)} = \beta. \quad (3.1)$$

*Proof.* Using (2.8), that is, the  $N = 2$  case of the Hardy-Ramanujan-Rademacher formula for  $p(n)$ , we find that

$$\log \sqrt[p(n)]{p(n)} = \frac{1}{n} \log \tilde{T}(n) + \frac{1}{n} \log(1 + \tilde{y}_n),$$

where  $\tilde{T}(n)$  and  $y_n$  are given by (2.9) and (2.13). By the definition (2.14) of  $\tilde{E}(n)$ , we get

$$\Delta^2 \log \sqrt[p(n-1)]{p(n-1)} = \Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) + \Delta^2 \tilde{E}(n-1). \quad (3.2)$$

Applying Lemma 2.2, we get that

$$\lim_{n \rightarrow +\infty} (n-1)^{\frac{5}{2}}\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1) = \beta. \quad (3.3)$$

From Lemma 2.3, we get

$$\lim_{n \rightarrow +\infty} (n-1)^{\frac{5}{2}}\Delta^2 \tilde{E}(n-1) = 0. \quad (3.4)$$

Using (3.2), (3.3) and (3.4), we deduce that

$$\lim_{n \rightarrow +\infty} n^{\frac{5}{2}}\Delta^2 \log \sqrt[p(n)]{p(n)} = \beta,$$

as required. ■

To prove Theorem 1.4, we need the following upper bound for  $\Delta^2 \log \sqrt[p(n-1)]{p(n-1)}$ .

**Theorem 3.2** *For  $n \geq 2$ ,*

$$\Delta^2 \log \sqrt[p(n-1)]{p(n-1)} < \frac{3\pi}{\sqrt{24}n^{5/2} + 3\pi}. \quad (3.5)$$

*Proof.* By the upper bound of  $\Delta^2 \frac{1}{n-1} \log \tilde{T}(n-1)$  given in Lemma 2.2, the upper bound of  $\Delta^2 \tilde{E}(n-1)$  given in Lemma 2.3, and the relation (3.2), we obtain the following upper bound of  $\Delta^2 \log \sqrt[n-1]{p(n-1)}$  for  $n \geq 40$ :

$$\Delta^2 \log \sqrt[n-1]{p(n-1)} < \frac{72\pi}{(n-1)(24n-25)^{3/2}} + \frac{5}{(n-1)^3} - \frac{4 \log[\mu(n+1)]}{(n+1)^3} + \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}.$$

To prove (3.5), we claim that for  $n \geq 2095$ ,

$$\frac{72\pi}{(n-1)(24n-25)^{3/2}} + \frac{5}{(n-1)^3} - \frac{4 \log[\mu(n+1)]}{(n+1)^3} + \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} < \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi}. \quad (3.6)$$

First, we show that for  $n \geq 60$ ,

$$\frac{72\pi}{(n-1)(24n-25)^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} < \frac{1}{(n-1)^3}. \quad (3.7)$$

For  $0 < x \leq \frac{1}{48}$ , it can be checked that

$$\frac{1}{(1-x)^{3/2}} < 1 + \frac{3}{2}x + \frac{3}{8}x^{\frac{3}{2}}. \quad (3.8)$$

In the notation  $\beta = 3\pi/\sqrt{24}$ , we have

$$\frac{72\pi}{(n-1)(24n-25)^{3/2}} = \frac{\beta}{(n-1)n^{3/2}(1-\frac{25}{24n})^{3/2}}. \quad (3.9)$$

Setting  $x = \frac{25}{24n}$ , we have  $x \leq \frac{1}{48}$  for  $n \geq 60$ . Applying (3.8) to the right hand side of (3.9), we find that for  $n \geq 60$ ,

$$\frac{\beta}{(n-1)n^{3/2}(1-\frac{25}{24n})^{3/2}} < \frac{\beta}{(n-1)n^{3/2}} \left[ 1 + \frac{75}{48n} + \frac{3}{8} \left( \frac{25}{24n} \right)^{\frac{3}{2}} \right], \quad (3.10)$$

so that for  $n \geq 60$ ,

$$\begin{aligned} & \frac{72\pi}{(n-1)[24n-25]^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} \\ & < \frac{\beta}{(n-1)n^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} + \frac{\beta}{(n-1)n^{3/2}} \left[ \frac{75}{48n} + \frac{3}{8} \left( \frac{25}{24n} \right)^{\frac{3}{2}} \right]. \end{aligned} \quad (3.11)$$

To prove (3.7), we proceed to show that the right hand side of (3.11) is bounded by  $\frac{1}{(n-1)^3}$ . Noting that for  $n \geq 2$ ,

$$\frac{\beta}{(n-1)n^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2}} + 3\pi} = \frac{\beta}{(n^{5/2} + \beta)(n-1)} + \frac{\beta^2}{(n^{5/2} + \beta)(n-1)n^{3/2}},$$

and using the fact  $n^{5/2} + \beta > (n-1)^{5/2}$ , together with  $n^{3/2} > (n-1)^{3/2}$ , we deduce that

$$\frac{\beta}{(n-1)n^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2} + 3\pi}} < \frac{\beta}{(n-1)^{7/2}} + \frac{\beta}{(n-1)^5}. \quad (3.12)$$

Applying (3.12) to (3.11), we obtain that for  $n \geq 60$ ,

$$\begin{aligned} & \frac{72\pi}{(n-1)[24n-25]^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2} + 3\pi}} \\ & < \frac{\beta}{(n-1)^{7/2}} + \frac{\beta^2}{(n-1)^5} + \frac{\beta}{(n-1)n^{3/2}} \left[ \frac{75}{48n} + \frac{3}{8} \left( \frac{25}{24n} \right)^{\frac{3}{2}} \right]. \end{aligned} \quad (3.13)$$

Since  $\frac{75}{48n} < \frac{2}{n-1}$  and  $\frac{3}{8} \left( \frac{25}{24n} \right)^{\frac{3}{2}} < \frac{1}{(n-1)^{3/2}}$  for  $n \geq 2$ , it follows from (3.13) that for  $n \geq 60$ ,

$$\begin{aligned} & \frac{72\pi}{(n-1)[24n-25]^{3/2}} - \frac{3\pi}{\sqrt{24n^{5/2} + 3\pi}} \\ & < \frac{\beta}{(n-1)^{7/2}} + \frac{\beta^2}{(n-1)^5} + \frac{2\beta}{(n-1)^{7/2}} + \frac{\beta}{(n-1)^4}. \end{aligned}$$

Using the fact that  $\beta < 2$ , we see that

$$\frac{3\beta}{(n-1)^{7/2}} + \frac{\beta^2}{(n-1)^5} + \frac{\beta}{(n-1)^4} < \frac{6}{(n-1)^{7/2}} + \frac{4}{(n-1)^5} + \frac{2}{(n-1)^4}. \quad (3.14)$$

For  $n \geq 60$ , it is easily checked that the right hand side of (3.14) is bounded by  $\frac{1}{(n-1)^3}$ . This confirms (3.7).

To prove the claim (3.6), it is enough to show that for  $n \geq 2095$ ,

$$\frac{1}{(n-1)^3} < \frac{4 \log[\mu(n+1)]}{(n+1)^3} - \frac{5}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}}. \quad (3.15)$$

From (2.61) it can be seen that for  $n \geq 2095$ ,

$$\frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} < \frac{5}{(n-1)^3}. \quad (3.16)$$

Since  $4 \log[\mu(n+1)] > 18$  for  $n \geq 2095$ , it follows from (3.16) that for  $n \geq 2095$ ,

$$\begin{aligned} & \frac{4 \log[\mu(n+1)]}{(n+1)^3} - \frac{5}{(n-1)^3} - \frac{5}{n-1} e^{-\frac{\pi\sqrt{24n-25}}{18}} \\ & > \frac{18}{(n+1)^3} - \frac{10}{(n-1)^3} > \frac{1}{(n-1)^3}. \end{aligned}$$

So we obtain (3.15). Combining (3.15) and (??), we arrive at (3.6). For  $2 \leq n \leq 2094$ , the inequality (3.5) can be easily checked. This completes the proof.  $\blacksquare$



We are now in a position to complete the proof of Theorem 1.4.

*Proof of Theorem 1.4.* It is known that for  $x > 0$ ,

$$\frac{x}{1+x} < \log(1+x),$$

so that for  $n \geq 1$ ,

$$\frac{3\pi}{\sqrt{24}n^{5/2} + 3\pi} < \log\left(1 + \frac{3\pi}{\sqrt{24}n^{5/2}}\right).$$

In light of the above relation, Theorem 3.2 implies that for  $n \geq 2$ ,

$$\Delta^2 \log \sqrt[n-1]{p(n-1)} < \log\left(1 + \frac{3\pi}{\sqrt{24}n^{5/2}}\right),$$

that is,

$$\sqrt[n+1]{p(n+1)} \sqrt[n-1]{p(n-1)} < \left(1 + \frac{3\pi}{\sqrt{24}n^{5/2}}\right) (\sqrt[n]{p(n)})^2,$$

as required. ■

We remark that  $\beta = 3\pi/\sqrt{24}$  is the smallest possible number for the inequality in Theorem 1.4. Suppose that  $0 < \gamma < \beta$ . By Theorem 3.1, there exists an integer  $N$  such that for  $n > N$ ,

$$n^{5/2} \Delta^2 \log \sqrt[n-1]{p(n-1)} > \gamma.$$

It follows that

$$\Delta^2 \log \sqrt[n-1]{p(n-1)} > \frac{\gamma}{n^{5/2}} > \log\left(1 + \frac{\gamma}{n^{5/2}}\right),$$

which implies that for  $n > N$ ,

$$\frac{\sqrt[n]{p(n)}}{\sqrt[n+1]{p(n+1)}} \left(1 + \frac{\gamma}{n^{5/2}}\right) < \frac{\sqrt[n-1]{p(n-1)}}{\sqrt[n]{p(n)}}.$$

**Acknowledgments.** We wish to thank the referee for helpful comments.

## References

- [1] G.E. Andrews, Combinatorial proof of a partition function limit, Amer. Math. Monthly., 78 (1971), 276–278.
- [2] G.E. Andrews, The Theory of Partitions, Cambridge University Press, Cambridge, 1998.
- [3] W.Y.C. Chen, J.J.F. Guo and L.X.W. Wang, Infinitely log-monotonic combinatorial sequences. Adv. Appl. Math., 52 (2014), 99–120.

- [4] W.Y.C. Chen, L.X.W. Wang and G.Y.B. Xie, Finite differences of the logarithm of the partition function. *Math. Comp.*, 85 (2016), 825–847.
- [5] S. Desalvo and I. Pak, Log-concavity of the partition function, *Ramanujan J.*, 38 (2016), 61–73.
- [6] G.H. Hardy, *Twelve Lectures on Subjects Suggested by His Life and Work*, Cambridge University Press, Cambridge, 1940.
- [7] G.H. Hardy and S. Ramanujan, Asymptotic formulae in combinatory analysis, *Proc. London Math. Soc.*, 17 (1918), 75–175.
- [8] D.H. Lehmer, On the series for the partition function, *Trans. Amer. Math. Soc.*, 43 (1938), 271–292.
- [9] D.H. Lehmer, On the remainders and convergence of the series for the partition function, *Trans. Amer. Math. Soc.*, 46 (1939), 362–373.
- [10] H. Rademacher, A convergent series for the partition function  $p(n)$ , *Proc. Nat. Acad. Sci.*, 23 (1937), 78–84.
- [11] Z.-W. Sun, On a sequence involving sums of primes, *Bull. Aust. Math. Soc.* 88 (2013), 197–205.
- [12] Y. Wang, B.-X. Zhu, Proofs of some conjectures on monotonicity of number-theoretic and combinatorial sequences, *Sci. China Math.* 57 (2014), 2429–2435.