




Contents lists available at ScienceDirect

Advances in Applied Mathematics

journal homepage: www.elsevier.com/locate/yaama

A unified framework to prove multiplicative inequalities for the partition function



Koustav Banerjee

Research Institute for Symbolic Computation, Johannes Kepler University,
Altenberger Strasse 69, A-4040 Linz, Austria

ARTICLE INFO

Article history:

Received 26 September 2022

Received in revised form 27 June 2023

Accepted 12 July 2023

Available online 28 July 2023

MSC:

05A16

05A20

11P82

Keywords:

Partition function

Hardy-Ramanujan-Rademacher formula

log-concavity

Finite difference

Partition inequalities

ABSTRACT

In this paper, we consider a certain class of inequalities for the partition function of the following form:

$$\prod_{i=1}^T p(n + s_i) \geq \prod_{i=1}^T p(n + r_i),$$

which we call multiplicative inequalities. Given a multiplicative inequality with the condition that $\sum_{i=1}^T s_i^m \neq \sum_{i=1}^T r_i^m$ for at least one $m \geq 1$, we shall construct a unified framework so as to decide whether such an inequality holds or not. As a consequence, we will see that study of such inequalities has manifold applications. For example, one can retrieve log-concavity property, strong log-concavity, and the multiplicative inequality for $p(n)$ considered by Bessenrodt and Ono, to name a few. Furthermore, we obtain an asymptotic expansion for the finite difference of the logarithm of $p(n)$, denoted by $(-1)^{r-1} \Delta^r \log p(n)$, which generalizes a result by Chen, Wang, and Xie.

© 2023 Elsevier Inc. All rights reserved.

E-mail address: Koustav.Banerjee@risc.uni-linz.ac.at.

<https://doi.org/10.1016/j.aam.2023.102590>

0196-8858/© 2023 Elsevier Inc. All rights reserved.

Contents

1. Introduction	2
2. Set up	5
3. Inequalities for $\log p(n; \vec{s})$	8
4. Asymptotics of $(-1)^{r-1} \Delta^r \log p(n)$	19
5. A framework to verify multiplicative inequalities for $p(n)$	25
6. Inequalities for $p(n; \vec{s})$	26
7. Conclusion	30
Acknowledgments	32
References	32

1. Introduction

A partition of a positive integer n is a weakly decreasing sequence $(\lambda_1, \lambda_2, \dots, \lambda_r)$ of positive integers such that $\lambda_1 + \lambda_2 + \dots + \lambda_r = n$. Let $p(n)$ denote the number of partitions of n . Hardy and Ramanujan [12] studied the asymptotic growth of $p(n)$ as follows:

$$p(n) \sim \frac{1}{4n\sqrt{3}} e^{\pi\sqrt{2n/3}} \text{ as } n \rightarrow \infty. \tag{1.1}$$

Rademacher [18,20,19] improved the work of Hardy and Ramanujan and found a convergent series for $p(n)$ and Lehmer’s [14,13] study was on estimation for the remainder term of the series for $p(n)$. The Hardy-Ramanujan-Rademacher formula reads

$$p(n) = \frac{\sqrt{12}}{24n-1} \sum_{k=1}^N \frac{A_k(n)}{\sqrt{k}} \left[\left(1 - \frac{k}{\mu(n)}\right) e^{\mu(n)/k} + \left(1 + \frac{k}{\mu(n)}\right) e^{-\mu(n)/k} \right] + R_2(n, N), \tag{1.2}$$

where

$$\mu(n) = \frac{\pi}{6} \sqrt{24n-1}, \quad A_k(n) = \sum_{\substack{h \pmod k \\ (h,k)=1}} e^{-2\pi i n h/k + \pi i s(h,k)}$$

with

$$s(h, k) = \sum_{\mu=1}^{k-1} \left(\frac{\mu}{k} - \left\lfloor \frac{\mu}{k} \right\rfloor - \frac{1}{2} \right) \left(\frac{h\mu}{k} - \left\lfloor \frac{h\mu}{k} \right\rfloor - \frac{1}{2} \right),$$

and

$$|R_2(n, N)| < \frac{\pi^2 N^{-2/3}}{\sqrt{3}} \left[\left(\frac{N}{\mu(n)} \right)^3 \sinh \frac{\mu(n)}{N} + \frac{1}{6} - \left(\frac{N}{\mu(n)} \right)^2 \right]. \tag{1.3}$$

A sequence $\{a_n\}_{n \geq 0}$ is said to satisfy the Turán inequalities or to be log-concave, if

$$a_n^2 - a_{n-1}a_{n+1} \geq 0 \text{ for all } n \geq 1. \tag{1.4}$$

Independently Nicolas [16] and DeSalvo and Pak [8, Theorem 1.1] proved that the partition function $p(n)$ is log-concave for all $n \geq 26$, conjectured by Chen [5]. DeSalvo and Pak [8, Theorem 4.1] also proved that for all $n \geq 2$,

$$\frac{p(n-1)}{p(n)} \left(1 + \frac{1}{n}\right) > \frac{p(n)}{p(n+1)}, \tag{1.5}$$

conjectured by Chen [5]. Further, they improved the rate of decay in (1.5) and proved that for all $n \geq 7$,

$$\frac{p(n-1)}{p(n)} \left(1 + \frac{240}{(24n)^{3/2}}\right) > \frac{p(n)}{p(n+1)}, \tag{1.6}$$

see [8, 4.2]. DeSalvo and Pak [8] finally came up with the conjecture that the coefficient of $1/n^{3/2}$ in (1.6) can be improved to $\pi/\sqrt{24}$; i.e., for all $n \geq 45$,

$$\frac{p(n-1)}{p(n)} \left(1 + \frac{\pi}{\sqrt{24}n^{3/2}}\right) > \frac{p(n)}{p(n+1)}, \tag{1.7}$$

which was proved by Chen, Wang and Xie [7, Sec. 2]. Recently, the author along with Paule, Radu, and Zeng [3, Theorem 7.6] confirmed that the coefficient of $1/n^{3/2}$ is indeed $\pi/\sqrt{24}$, which is the optimal; i.e., they proved that for all $n \geq 120$,

$$p(n)^2 > \left(1 + \frac{\pi}{\sqrt{24}n^{3/2}} - \frac{1}{n^2}\right)p(n-1)p(n+1). \tag{1.8}$$

DeSalvo and Pak [8, Theorem 5.1] also established that $p(n)$ satisfies the strong log-concavity property; i.e., for all $n > m > 1$,

$$p(n)^2 - p(n-m)p(n+m) > 0. \tag{1.9}$$

Ono and Bessenrodt [4] extended (1.9) by considering the border case $m = n$. This leads to unveil multiplicative properties of the partition function encoded in the following theorem.

Theorem 1.1. [4, Theorem 2.1] *If a and b are integers with $a, b > 1$ and $a + b > 8$, then*

$$p(a)p(b) \geq p(a+b), \tag{1.10}$$

with equality holding only for $\{a, b\} = \{2, 7\}$.

Let Δ be the forward difference operator define by $\Delta a(n) := a(n+1) - a(n)$ for a sequence $(a(n))_{n \geq 0}$. It is clear that the log-concavity property for $p(n)$ is equivalent to say that $-\Delta^2 \log p(n-1) > 0$ for all $n \geq 26$. Equations (1.7) and (1.8) show the asymptotic growth of $-\Delta^2 \log p(n-1)$. Chen, Wang, and Xie proved the positivity of $(-1)^{r-1} \Delta^r \log p(n)$ along with estimation of an upper bound.

Theorem 1.2. [7, Thm. 3.1 and 4.1] For each $r \geq 1$, there exists a positive integer $n(r)$ such that for all $n \geq n(r)$,

$$0 < (-1)^{r-1} \Delta^r \log p(n) < \log \left(1 + \frac{\pi}{\sqrt{6}} \left(\frac{1}{2} \right)_{r-1} \frac{1}{(n+1)^{r-\frac{1}{2}}} \right). \quad (1.11)$$

The above inequalities can be rephrased in the following form:

$$\prod_{i=1}^T p(n+s_i) \geq \prod_{i=1}^T p(n+r_i), \quad (1.12)$$

which we call multiplicative inequalities for the partition function. Instead of applying the Hardy-Ramanujan-Rademacher formula (1.2) and Lehmer's error bound (1.3) but with different methodology for different inequalities for $p(n)$ as done in [4,7,8,16], we will see how one can prove all such multiplicative inequalities under a unified framework so as to decide explicitly $N(T)$, such that for all $n \geq N(T)$, (1.12) holds. To prove (1.12), it is equivalent to show

$$\sum_{i=1}^T \log p(n+s_i) \geq \sum_{i=1}^T \log p(n+r_i), \quad (1.13)$$

and therefore, an infinite family of inequalities for logarithm of the shifted version of the partition function is a prerequisite, see Theorems 3.9 and 3.13. As an application of Theorem 3.9, we shall complete Theorem 1.2 (see Theorems 4.9 and 4.10 below) in the following aspects:

- (1) by improving the lower bound in (1.11) to show that the rate of decay given in the upper bound is the optimal one,
- (2) for each $r \geq 1$, computation of $n(r)$ by estimating error bound based on the minimal choice of the truncation point w in Theorem 3.9,
- (3) and a full asymptotic expansion for $(-1)^{r-1} \Delta^r \log p(n)$. This seems to be inaccessible from Theorem 1.2 because a key tool in the proof was on the relations between the higher order differences and derivatives (cf. Prop. 3.5, [7]) due to Odlyzko [17] which only contributes to the main term in the expansion; i.e., $\frac{\pi}{\sqrt{6}} \left(\frac{1}{2} \right)_{r-1} \frac{1}{(n+1)^{r-\frac{1}{2}}}$.

Even having Theorem 3.13 in hand, in order to decide whether (1.12) holds or not, there are two key factors remain unexplained. First, an explanation of the following assumption

$$\sum_{i=1}^T s_i^m \neq \sum_{i=1}^T r_i^m \text{ for at least one } m \geq \mathbb{Z}_{\geq 1}, \tag{1.14}$$

and an appropriate choice of w , i.e., the truncation point as in Theorem 3.13. Now we move on to see how these two factors are intricately connected through a classical problem in Diophantine equations known as the Prouhet-Tarry-Escott problem [9, Chapter XXIV]. The problem originated in different guise from a letter of Goldbach [10] to Euler that dates back to 18 July, 1750. The Prouhet-Tarry-Escott problem asks for two distinct tuples of integers (s_1, s_2, \dots, s_T) and (r_1, r_2, \dots, r_T) such that

$$\sum_{i=1}^T s_i^k = \sum_{i=1}^T r_i^k, \text{ for all } 0 \leq k \leq m - 1 \text{ and } \sum_{i=1}^T s_i^m \neq \sum_{i=1}^T r_i^m.$$

We write $(s_1, \dots, s_T) \stackrel{m}{=} (r_1, \dots, r_T)$ to denote a solution of the Prouhet-Tarry-Escott problem. Recently, Merca and Katriel [15] connects the non-trivial linear homogeneous partition inequalities with the Prouhet-Tarry-Escott problem. In brevity, we shall explain why the optimal choice of truncation point $w = m + 1$, with $(s_1, \dots, s_T) \stackrel{m}{=} (r_1, \dots, r_T)$ for a given (1.12) in Section 5.

The rest of the paper is organized as follows. In Section 2, we state preliminary lemmas and theorems from the work of Paule, Radu, Zeng, and the author [3]. Section 3 presents a detailed synthesis on derivation of inequalities for $\log p(n+s)$ for any non-negative integer s that leads to the main result of this paper, see Theorem 3.13. As an application of Theorem 3.13, we provide a full asymptotic expansion of $(-1)^{r-1} \Delta^r \log p(n)$ in Section 4. In Section 5, we work out the steps to verify multiplicative inequalities for the partition function. Section 6 is devoted to derive an infinite family of inequalities for $\prod_{i=1}^T p(n + s_i)$, given in Theorem 6.9. Finally we conclude this paper with a short discussion on the applications of Theorems 3.13 and 6.9.

2. Set up

Throughout this section, we follow the notations as in [3].

Definition 2.1 (Def. 5.1, [3]). For $y \in \mathbb{R}$, $0 < y^2 < 24$, we define

$$G(y) := -\log\left(1 - \frac{y^2}{24}\right) + \frac{\pi\sqrt{24}}{6y} \left(\sqrt{1 - \frac{y^2}{24}} - 1\right) + \log\left(1 - \frac{y}{\frac{\pi}{6}\sqrt{24 - y^2}}\right), \tag{2.1}$$

and its sequence of Taylor coefficients by

$$G(y) = \sum_{u=1}^{\infty} g_u y^u. \tag{2.2}$$

Define $\alpha := \frac{\pi^2}{36 + \pi^2}$.

Lemma 2.2 (*Lem. 5.4, [3]*). Let $G(y) = \sum_{u=1}^{\infty} g_u y^u$ be the Taylor expansion of $G(y)$ as in Definition 2.1. Then for $n \geq 1$,

$$g_{2n} = \frac{1}{3^n 2^{3n} n} - \frac{1}{3^n 2^{3n+1} n} \left(-1 + \frac{1}{\alpha^n}\right), \tag{2.3}$$

and for $n \geq 0$,

$$g_{2n+1} = \sqrt{6} \left[(-1)^{n+1} \binom{1/2}{n+1} \frac{\pi}{2^{3n+3} 3^{n+2}} - \frac{1}{2^{3n+1} 3^n \alpha^n (2n+1) \pi} \sum_{j=0}^n \alpha^j \binom{-\frac{1}{2} + j}{j} \right]. \tag{2.4}$$

Lemma 2.3 (*Lem. 5.8, [3]*). For $n \geq 0$, we have

$$-\frac{\sqrt{6}}{2\pi 2^{3n} 3^n \alpha^n (2n+1)} \left(\frac{\pi^2}{72} + 1 + \frac{\alpha}{2(1-\alpha)} \right) \leq g_{2n+1} \leq -\frac{\sqrt{6}}{2\pi 2^{3n} 3^n \alpha^n (2n+1)} \left(1 + \frac{\alpha}{2} \right). \tag{2.5}$$

Lemma 2.4 (*Lem. 5.9, [3]*). For $n \geq 1$, we have

$$-\frac{1}{3^n 2^{3n+1} \alpha^n n} \leq g_{2n} \leq \frac{1}{3^n 2^{3n} \alpha^n n} \left(\frac{3\alpha}{2} - \frac{1}{2} \right). \tag{2.6}$$

Definition 2.5 (*Def. 4.3, [3]*). For $k \in \mathbb{Z}_{\geq 2}$, define

$$g(k) := \frac{1}{24} \left(\frac{6^2}{\pi^2} \cdot \nu(k)^2 + 1 \right),$$

where $\nu(k) := 2 \log 6 + (2 \log 2)k + 2k \log k + 2k \log \log k + \frac{5k \log \log k}{\log k}$.

Definition 2.6 (*Def. 6.4, [3]*). For $n, U \in \mathbb{Z}_{\geq 1}$, we define

$$P_n(U) := -\log 4\sqrt{3} - \log n + \pi \sqrt{\frac{2n}{3}} + \sum_{u=1}^U g_u (1/\sqrt{n})^u.$$

Theorem 2.7 (Thm 6.6, [3]). Let $G(y) = \sum_{u=1}^{\infty} g_u y^u$ as in Definition 2.1. Let $g(k)$ be as in Definition 2.5 and $P_n(U)$ as in Definition 2.6. If $m \geq 1$ and $n > g(2m)$, then

$$P_n(2m - 1) - \frac{2}{3^m 2^{3m} \alpha^m n^m} < \log p(n) < P_n(2m - 1) + \frac{1}{3^m 2^{3m} \alpha^m n^m}; \quad (2.7)$$

if $m \geq 2$ and $n > g(2m - 1)$, then

$$P_n(2m - 2) - \frac{7}{3^m 2^{3m} \alpha^m n^{m-1/2} (2m - 1)} < \log p(n) < P_n(2m - 2) + \frac{2}{3^m 2^{3m} \alpha^m n^{m-1/2} (2m - 1)}. \quad (2.8)$$

In other words, for $w \in \mathbb{Z}_{>0}$ with $\lceil w/2 \rceil \geq \gamma_0$ and $n > g(w)$, we have

$$P_n(w - 1) - \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil}} \left(\frac{1}{\sqrt{n}}\right)^w < \log p(n) < P_n(w - 1) + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil}} \left(\frac{1}{\sqrt{n}}\right)^w, \quad (2.9)$$

where

$$(\gamma_0, \gamma_1, \gamma_2) = \begin{cases} (1, 4, 2), & \text{if } w \text{ is even} \\ (2, 7, 2), & \text{if } w \text{ is odd} \end{cases}. \quad (2.10)$$

Lemma 2.8 (Lem 7.3, [3]). For $n, s \in \mathbb{Z}_{\geq 1}, m \in \mathbb{N}$ and $n > 2s$, let

$$b_{m,n}(s) := \frac{4\sqrt{s}}{\sqrt{s+m-1}} \binom{s+m-1}{s-1} \frac{1}{n^m},$$

then

$$-b_{m,n}(s) < \sum_{k=m}^{\infty} \binom{-\frac{2s-1}{2}}{k} \frac{1}{n^k} < b_{m,n}(s) \quad (2.11)$$

and

$$0 < \sum_{k=m}^{\infty} \binom{-\frac{2s-1}{2}}{k} \frac{(-1)^k}{n^k} < b_{m,n}(s). \quad (2.12)$$

Lemma 2.9 (Lem 7.4, [3]). For $n, s \in \mathbb{Z}_{\geq 1}, m \in \mathbb{N}$ and $n > 2s$, let

$$\beta_{m,n}(s) := \frac{2}{n^m} \binom{s+m-1}{s-1},$$

then

$$-\beta_{m,n}(s) < \sum_{k=m}^{\infty} \binom{-s}{k} \frac{1}{n^k} < \beta_{m,n}(s) \tag{2.13}$$

and

$$0 < \sum_{k=m}^{\infty} \binom{-s}{k} \frac{(-1)^k}{n^k} < \beta_{m,n}(s). \tag{2.14}$$

Lemma 2.10 (*Lem 7.5, [3]*). For $m, n, s \in \mathbb{Z}_{\geq 1}$ and $n > 2s$, let

$$c_{m,n}(s) := \frac{2}{m} \frac{s^m}{n^m},$$

then

$$-c_{m,n}(s) < \sum_{k=m}^{\infty} \frac{(-1)^{k+1} s^k}{k n^k} < c_{m,n}(s) \quad \text{and} \quad -c_{m,n}(s) < -\sum_{k=m}^{\infty} \frac{1}{k} \frac{s^k}{n^k} < 0 \tag{2.15}$$

and

$$-\frac{c_{m,n}(s)}{\sqrt{m}} < \sum_{k=m}^{\infty} \binom{1/2}{k} \frac{s^k}{n^k} < \frac{c_{m,n}(s)}{\sqrt{m}} \quad \text{and} \quad -\frac{c_{m,n}(s)}{\sqrt{m}} < \sum_{k=m}^{\infty} \binom{1/2}{k} \frac{(-1)^k s^k}{n^k} < 0. \tag{2.16}$$

3. Inequalities for $\log p(n; \vec{s})$

In this section, first we prove an infinite family of inequalities for $\log p(n + s)$ with s being a non-negative integer, see Theorem 3.9. Starting from Theorem 2.7, we will estimate $P_{n+s}(U)$ and the error terms given in (2.7) and (2.8), stated in Lemma 3.3-3.6. Finally, generalizing Theorem 3.9 by taking into consideration $\sum_{i=1}^T \log p(n + s_i)$ for $(s_1, s_2, \dots, s_T) \in \mathbb{Z}_{\geq 0}^T$, we obtain Theorem 3.13.

Lemma 3.1. *Let the coefficient sequence $(g_n)_{n \geq 1}$ be as in Lemma 2.2. Then for all $n \geq 1$, we have*

$$|g_n| \leq \frac{1}{n} \frac{1}{(24\alpha)^{\lfloor n/2 \rfloor}}. \tag{3.1}$$

Proof. Observe that for all $n \geq 0$, $\frac{\sqrt{6}}{2\pi} \left(1 + \frac{\alpha}{2}\right) \frac{1}{(24\alpha)^n (2n+1)} > 0$ and $0 < \frac{\sqrt{6}}{2\pi} \left(\frac{\pi^2}{72} + 1 + \frac{\alpha}{2(1-\alpha)}\right) < 1$. Using (2.5), we obtain for all $n \geq 0$,

$$-\frac{1}{(24\alpha)^n (2n+1)} < g_{2n+1} < 0. \tag{3.2}$$

Since $\frac{3\alpha}{2} - \frac{1}{2} < 0$, from (2.6), it follows that for all $n \geq 1$,

$$-\frac{1}{(24\alpha)^n(2n)} \leq g_{2n} < 0. \tag{3.3}$$

From (3.2) and (3.3), we conclude that for all $n \geq 1$,

$$|g_n| \leq \frac{1}{(24\alpha)^{\lfloor n/2 \rfloor} n}. \quad \square$$

Definition 3.2. For $s \in \mathbb{Z}_{\geq 0}$, define

$$\delta_s := \begin{cases} 1, & \text{if } s \geq 1 \\ 0, & \text{if } s = 0 \end{cases}.$$

Lemma 3.3. For $(n, s) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 0}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s$, let

$$P_{n,s}^1(w) := -\log n + \sum_{k=1}^{\lfloor \frac{w-1}{2} \rfloor} \frac{(-1)^k s^k}{k} \left(\frac{1}{\sqrt{n}}\right)^{2k} \quad \text{and} \quad E_{n,s}^1(w) := \frac{2s^{\lceil \frac{w+1}{2} \rceil}}{\lceil w/2 \rceil} \left(\frac{1}{\sqrt{n}}\right)^w \delta_s,$$

then

$$P_{n,s}^1(w) - E_{n,s}^1(w) \leq -\log(n+s) \leq P_{n,s}^1(w) + E_{n,s}^1(w). \tag{3.4}$$

Proof. For all $n, s \in \mathbb{Z}_{\geq 1}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s$, we split $\log(n+s)$ as follows

$$\log(n+s) = \log n + \sum_{k=1}^{\infty} \frac{(-1)^{k+1} s^k}{k} \frac{1}{n^k} = \log n + \sum_{k=1}^{\lfloor \frac{w-1}{2} \rfloor} \frac{(-1)^{k+1} s^k}{k} \frac{1}{n^k} + \sum_{k=\lceil \frac{w}{2} \rceil}^{\infty} \frac{(-1)^{k+1} s^k}{k} \frac{1}{n^k}. \tag{3.5}$$

Applying (2.15) with $m \mapsto \lceil \frac{w}{2} \rceil$, it follows that for all $n > 2s$,

$$-\frac{2}{\lceil w/2 \rceil} \left(\frac{s}{n}\right)^{\lceil w/2 \rceil} < \sum_{k=\lceil \frac{w}{2} \rceil}^{\infty} \frac{(-1)^{k+1} s^k}{k} \frac{1}{n^k} < \frac{2}{\lceil w/2 \rceil} \left(\frac{s}{n}\right)^{\lceil w/2 \rceil}. \tag{3.6}$$

Since for all $s \in \mathbb{Z}_{\geq 0}$, $s^{\lceil w/2 \rceil} \leq s^{\lceil \frac{w+1}{2} \rceil}$, from (3.5) and (3.6), it follows that

$$P_{n,s}^1(w) - E_{n,s}^1(w) \leq -\log(n+s) \leq P_{n,s}^1(w) + E_{n,s}^1(w). \tag{3.7}$$

Observe that equality holds when $s = 0$. \square

Lemma 3.4. For $(n, s) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 0}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s$, let

$$P_{n,s}^2(w) := \pi \sqrt{\frac{2n}{3}} + \pi \sqrt{\frac{2}{3}} \sum_{k=1}^{\lfloor \frac{w}{2} \rfloor} \binom{1/2}{k} s^k \left(\frac{1}{\sqrt{n}}\right)^{2k-1} \quad \text{and} \quad E_{n,s}^2(w) := \frac{6s^{\lceil \frac{w+1}{2} \rceil}}{\lceil w/2 \rceil} \left(\frac{1}{\sqrt{n}}\right)^w \delta_s,$$

then

$$P_{n,s}^2(w) - E_{n,s}^2(w) \leq \pi \sqrt{\frac{2n+2s}{3}} \leq P_{n,s}^2(w) + E_{n,s}^2(w). \tag{3.8}$$

Proof. For all $n, s \in \mathbb{Z}_{\geq 1}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s$, we split $\pi \sqrt{\frac{2n+2s}{3}}$ as follows

$$\pi \sqrt{\frac{2n+2s}{3}} = \pi \sqrt{\frac{2n}{3}} + \pi \sqrt{\frac{2}{3}} \sum_{k=1}^{\lfloor \frac{w}{2} \rfloor} \binom{1/2}{k} s^k \left(\frac{1}{\sqrt{n}}\right)^{2k-1} + \pi \sqrt{\frac{2n}{3}} \sum_{k=\lfloor \frac{w+2}{2} \rfloor}^{\infty} \binom{1/2}{k} \frac{s^k}{n^k}. \tag{3.9}$$

Applying (2.16) with $m \mapsto \lfloor \frac{w+2}{2} \rfloor$, it follows that for all $n > 2s$,

$$-\frac{2}{\left(\lfloor \frac{w+2}{2} \rfloor\right)^{3/2}} \left(\frac{s}{n}\right)^{\lfloor \frac{w+2}{2} \rfloor} < \sum_{k=\lfloor \frac{w+2}{2} \rfloor}^{\infty} \binom{1/2}{k} \frac{s^k}{n^k} < \frac{2}{\left(\lfloor \frac{w+2}{2} \rfloor\right)^{3/2}} \left(\frac{s}{n}\right)^{\lfloor \frac{w+2}{2} \rfloor}. \tag{3.10}$$

Therefore,

$$\begin{aligned} -2\pi \sqrt{\frac{2}{3}} \frac{s^{\lfloor \frac{w+2}{2} \rfloor}}{\left(\lfloor \frac{w+2}{2} \rfloor\right)^{3/2}} \left(\frac{1}{\sqrt{n}}\right)^{2\lfloor \frac{w+2}{2} \rfloor - 1} &< \pi \sqrt{\frac{2n}{3}} \sum_{k=\lfloor \frac{w+2}{2} \rfloor}^{\infty} \binom{1/2}{k} \frac{s^k}{n^k} \\ &< 2\pi \sqrt{\frac{2}{3}} \frac{s^{\lfloor \frac{w+2}{2} \rfloor}}{\left(\lfloor \frac{w+2}{2} \rfloor\right)^{3/2}} \left(\frac{1}{\sqrt{n}}\right)^{2\lfloor \frac{w+2}{2} \rfloor - 1}. \end{aligned} \tag{3.11}$$

Now for all $s \in \mathbb{Z}_{\geq 0}$,

$$\pi \sqrt{\frac{2}{3}} \frac{s^{\lfloor \frac{w+2}{2} \rfloor}}{\left(\lfloor \frac{w+2}{2} \rfloor\right)^{3/2}} \left(\frac{1}{\sqrt{n}}\right)^{2\lfloor \frac{w+2}{2} \rfloor - 1} < \frac{6s^{\lceil \frac{w+1}{2} \rceil}}{\lceil w/2 \rceil} \left(\frac{1}{\sqrt{n}}\right)^w.$$

From (3.9) and (3.11), it follows that

$$P_{n,s}^2(w) - E_{n,s}^2(w) \leq \pi \sqrt{\frac{2n+2s}{3}} \leq P_{n,s}^2(w) + E_{n,s}^2(w), \tag{3.12}$$

with equality holds for $s = 0$. \square

Lemma 3.5. For $(n, s) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 0}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s$, let

$$\begin{aligned} \bar{g}_\ell(s; t) &:= g_\ell \left(\begin{matrix} -\ell/2 \\ t - \lfloor \ell/2 \rfloor \end{matrix} \right) s^{t - \lfloor \frac{\ell}{2} \rfloor} \quad \text{for all } \ell \in \mathbb{Z}_{\geq 1}, \\ P_{n,s}^3(w) &:= \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n}} \right)^u + \sum_{t=1}^{\lfloor \frac{w-2}{2} \rfloor} \sum_{u=0}^{t-1} \bar{g}_{2u+1}(s; t) \left(\frac{1}{\sqrt{n}} \right)^{2t+1} \\ &\quad + \sum_{t=2}^{\lfloor \frac{w-1}{2} \rfloor} \sum_{u=1}^{t-1} \bar{g}_{2u}(s; t) \left(\frac{1}{\sqrt{n}} \right)^{2t}, \end{aligned}$$

and

$$E_{n,s}^3(w) := \frac{29}{w} \left(s + \frac{1}{24\alpha} \right)^{\lceil \frac{w-1}{2} \rceil + 1} \left(\frac{1}{\sqrt{n}} \right)^w \delta_s,$$

then

$$P_{n,s}^3(w) - E_{n,s}^3(w) \leq \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n+s}} \right)^u \leq P_{n,s}^3(w) + E_{n,s}^3(w). \tag{3.13}$$

Proof. For all $n, s \in \mathbb{Z}_{\geq 1}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s$, we split $\sum_{u=1}^{w-1} g_u(1/\sqrt{n+s})^u$ as

$$\begin{aligned} \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n+s}} \right)^u &= \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n}} \right)^u \sum_{k=0}^{\infty} \binom{-u/2}{k} \frac{s^k}{n^k} \\ &= \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n}} \right)^u + \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n}} \right)^u \sum_{k=1}^{\infty} \binom{-u/2}{k} \frac{s^k}{n^k} \\ &= \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n}} \right)^u + \sum_{u=1}^{w-1} g_u \sum_{k=1}^{\infty} \binom{-u/2}{k} s^k \left(\frac{1}{\sqrt{n}} \right)^{2k+u} \\ &= \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n}} \right)^u + \sum_{u=0}^{\lfloor \frac{w-2}{2} \rfloor} g_{2u+1} \sum_{k=1}^{\infty} \binom{-2u+1}{k} s^k \left(\frac{1}{\sqrt{n}} \right)^{2k+2u+1} \\ &\quad + \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} g_{2u} \sum_{k=1}^{\infty} \binom{-u}{k} s^k \left(\frac{1}{\sqrt{n}} \right)^{2k+2u}. \tag{3.14} \end{aligned}$$

Now,

$$\sum_{u=0}^{\lfloor \frac{w-2}{2} \rfloor} g_{2u+1} \sum_{k=1}^{\infty} \binom{-2u+1}{k} s^k \left(\frac{1}{\sqrt{n}} \right)^{2k+2u+1}$$

$$\begin{aligned}
 &= \sum_{u=0}^{\lfloor \frac{w-2}{2} \rfloor} g_{2u+1} \sum_{t=u+1}^{\infty} \binom{-2u+1}{t-u} s^{t-u} \left(\frac{1}{\sqrt{n}}\right)^{2t+1} \\
 &= \sum_{u=0}^{\lfloor \frac{w-2}{2} \rfloor} g_{2u+1} \sum_{t=u+1}^{\lfloor \frac{w-2}{2} \rfloor} \binom{-2u+1}{t-u} s^{t-u} \left(\frac{1}{\sqrt{n}}\right)^{2t+1} \\
 &\quad + \sum_{u=0}^{\lfloor \frac{w-2}{2} \rfloor} g_{2u+1} \sum_{t=\lceil \frac{w-1}{2} \rceil}^{\infty} \binom{-2u+1}{t-u} s^{t-u} \left(\frac{1}{\sqrt{n}}\right)^{2t+1} \\
 &= \sum_{t=1}^{\lfloor \frac{w-2}{2} \rfloor} \sum_{u=0}^{t-1} \bar{g}_{2u+1}(s; t) \left(\frac{1}{\sqrt{n}}\right)^{2t+1} \\
 &\quad + \underbrace{\sum_{u=1}^{\lceil \frac{w-1}{2} \rceil} \sum_{t=\lceil \frac{w-1}{2} \rceil - u + 1}^{\infty} g_{2u-1} \binom{-2u-1}{t} s^t \left(\frac{1}{\sqrt{n}}\right)^{2t+2u-1}}_{=: \mathcal{S}_o(w, n, s)}. \tag{3.15}
 \end{aligned}$$

Next, we proceed to estimate the absolute value of the error sum $\mathcal{S}_o(w, n, s)$ for $s \in \mathbb{Z}_{\geq 1}$.

$$\begin{aligned}
 &|\mathcal{S}_o(w, n, s)| \\
 &\leq \sum_{u=1}^{\lceil \frac{w-1}{2} \rceil} |g_{2u-1}| \left(\frac{1}{\sqrt{n}}\right)^{2u-1} \left| \sum_{t=\lceil \frac{w-1}{2} \rceil - u + 1}^{\infty} \binom{-2u-1}{t} \frac{s^t}{n^t} \right| \\
 &< 4 \sum_{u=1}^{\lceil \frac{w-1}{2} \rceil} |g_{2u-1}| \left(\frac{1}{\sqrt{n}}\right)^{2u-1} \sqrt{\frac{u}{\lceil \frac{w-1}{2} \rceil}} \binom{\lceil \frac{w-1}{2} \rceil}{u-1} \left(\frac{s}{n}\right)^{\lceil \frac{w-1}{2} \rceil - u + 1} \\
 &\quad \left(\text{by substitution } (m, s, n) \mapsto \left(\lceil \frac{w-1}{2} \rceil - u + 1, u, \frac{n}{s}\right) \text{ in (2.11)} \right) \\
 &\leq 4 \left(\sum_{u=1}^{\lceil \frac{w-1}{2} \rceil} |g_{2u-1}| \binom{\lceil \frac{w-1}{2} \rceil}{u-1} \frac{1}{s^u} \right) \left(\frac{1}{\sqrt{n}}\right)^{2\lceil \frac{w-1}{2} \rceil + 1} s^{\lceil \frac{w-1}{2} \rceil + 1} \\
 &\leq 4 \left(\sum_{u=1}^{\lceil \frac{w-1}{2} \rceil} \frac{1}{(2u-1)(24\alpha)^{u-1}} \binom{\lceil \frac{w-1}{2} \rceil}{u-1} \frac{1}{s^u} \right) \left(\frac{1}{\sqrt{n}}\right)^w s^{\lceil \frac{w-1}{2} \rceil + 1} \quad (\text{by Lemma 3.1}) \\
 &= 4 \left(\sum_{u=0}^{\lceil \frac{w-1}{2} \rceil - 1} \frac{1}{2u+1} \binom{\lceil \frac{w-1}{2} \rceil}{u} \frac{1}{(24\alpha s)^u} \right) \left(\frac{1}{\sqrt{n}}\right)^w s^{\lceil \frac{w-1}{2} \rceil} \\
 &\leq \frac{16}{3} \left(\sum_{u=0}^{\lceil \frac{w-1}{2} \rceil - 1} \frac{1}{2u+2} \binom{\lceil \frac{w-1}{2} \rceil}{u} \frac{1}{(24\alpha s)^u} \right) \left(\frac{1}{\sqrt{n}}\right)^w s^{\lceil \frac{w-1}{2} \rceil}
 \end{aligned}$$

$$\begin{aligned}
 & \left(\text{as } \frac{4}{2u+1} \leq \frac{8}{3(u+1)} \text{ for all } u \geq 1 \right) \\
 &= \frac{16}{3} \frac{1}{2^{\lceil \frac{w-1}{2} \rceil + 1}} \left(\left(\left(1 + \frac{1}{24\alpha s} \right)^{\lceil \frac{w-1}{2} \rceil + 1} - 1 \right) 24\alpha s - \left(\frac{1}{24\alpha s} \right)^{\lceil \frac{w-1}{2} \rceil} \right) \\
 & \quad \times \left(\frac{1}{\sqrt{n}} \right)^w s^{\lceil \frac{w-1}{2} \rceil} \\
 &< \frac{16}{3w} \left(\left(1 + \frac{1}{24\alpha s} \right)^{\lceil \frac{w-1}{2} \rceil + 1} - 1 \right) 24\alpha s \left(\frac{1}{\sqrt{n}} \right)^w s^{\lceil \frac{w-1}{2} \rceil} \\
 &< \frac{28}{w} \left(s + \frac{1}{24\alpha} \right)^{\lceil \frac{w-1}{2} \rceil + 1} \left(\frac{1}{\sqrt{n}} \right)^w. \tag{3.16}
 \end{aligned}$$

Similar to (3.15), we get

$$\begin{aligned}
 & \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} g_{2u} \sum_{k=1}^{\infty} \binom{-u}{k} s^k \left(\frac{1}{\sqrt{n}} \right)^{2k+2u} \\
 &= \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} \sum_{t=u+1}^{\infty} g_{2u} \binom{-u}{t-u} s^{t-u} \left(\frac{1}{\sqrt{n}} \right)^{2t} \\
 &= \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} \sum_{t=u+1}^{\lfloor \frac{w-1}{2} \rfloor} g_{2u} \binom{-u}{t-u} s^{t-u} \left(\frac{1}{\sqrt{n}} \right)^{2t} + \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} \sum_{t=\lceil \frac{w}{2} \rceil}^{\infty} g_{2u} \binom{-u}{t-u} s^{t-u} \left(\frac{1}{\sqrt{n}} \right)^{2t} \\
 &= \sum_{t=2}^{\lfloor \frac{w-1}{2} \rfloor} \sum_{u=1}^{t-1} \bar{g}_{2u}(s; t) \left(\frac{1}{\sqrt{n}} \right)^{2t} + \underbrace{\sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} \sum_{t=\lceil \frac{w}{2} \rceil - u}^{\infty} g_{2u} \binom{-u}{t} s^t \left(\frac{1}{\sqrt{n}} \right)^{2t+2u}}_{=: \mathcal{S}_e(w, n, s)}. \tag{3.17}
 \end{aligned}$$

Consequently for $s \in \mathbb{Z}_{\geq 1}$,

$$\begin{aligned}
 |\mathcal{S}_e(w, n, s)| &\leq \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} |g_{2u}| \left(\frac{1}{\sqrt{n}} \right)^{2u} \left| \sum_{t=\lceil \frac{w}{2} \rceil - u}^{\infty} \binom{-u}{t} \frac{s^t}{n^t} \right| \\
 &< 2 \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} |g_{2u}| \left(\frac{1}{\sqrt{n}} \right)^{2u} \binom{\lceil \frac{w}{2} \rceil - 1}{u-1} \left(\frac{s}{n} \right)^{\lceil \frac{w}{2} \rceil - u} \\
 & \quad \left(\text{by substitution } (m, s, n) \mapsto \left(\lceil \frac{w}{2} \rceil - u, u, \frac{n}{s} \right) \text{ in (2.13)} \right) \\
 &= 2 \left(\sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} |g_{2u}| \binom{\lceil \frac{w}{2} \rceil - 1}{u-1} \frac{1}{s^u} \right) s^{\lceil \frac{w}{2} \rceil} \left(\frac{1}{\sqrt{n}} \right)^w
 \end{aligned}$$

$$\begin{aligned}
 &\leq 2 \left(\sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} \frac{1}{2u} \binom{\lceil \frac{w}{2} \rceil - 1}{u-1} \frac{1}{(24\alpha s)^u} \right) s^{\lceil \frac{w}{2} \rceil} \left(\frac{1}{\sqrt{n}} \right)^w \quad (\text{by Lemma 3.1}) \\
 &= 2 \left(\sum_{u=1}^{\lceil \frac{w}{2} \rceil - 1} \frac{1}{2u} \binom{\lceil \frac{w}{2} \rceil - 1}{u-1} \frac{1}{(24\alpha s)^u} \right) s^{\lceil \frac{w}{2} \rceil} \left(\frac{1}{\sqrt{n}} \right)^w \\
 &= \frac{1}{w} \left(\left(1 + \frac{1}{24\alpha s} \right)^{\lceil \frac{w}{2} \rceil} - 1 - \left(\frac{1}{24\alpha s} \right)^{\lceil \frac{w}{2} \rceil} \right) s^{\lceil \frac{w}{2} \rceil} \left(\frac{1}{\sqrt{n}} \right)^w \\
 &< \frac{1}{w} \left(s + \frac{1}{24\alpha} \right)^{\lceil \frac{w-1}{2} \rceil + 1} \left(\frac{1}{\sqrt{n}} \right)^w. \tag{3.18}
 \end{aligned}$$

From (3.14), (3.15), and (3.17), we obtain

$$\sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n+s}} \right)^u - P_{n,s}^3(w) = \mathcal{S}_o(w, n, s) + \mathcal{S}_e(w, n, s), \tag{3.19}$$

and taking absolute on both sides of (3.19) and applying (3.16) and (3.18), it follows that

$$\begin{aligned}
 \left| \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n+s}} \right)^u - P_{n,s}^3(w) \right| &= \left| \mathcal{S}_o(w, n, s) + \mathcal{S}_e(w, n, s) \right| \\
 &\leq \left| \mathcal{S}_o(w, n, s) \right| + \left| \mathcal{S}_e(w, n, s) \right| \tag{3.20} \\
 &< \frac{29}{w} \left(s + \frac{1}{24\alpha} \right)^{\lceil \frac{w-1}{2} \rceil + 1} \left(\frac{1}{\sqrt{n}} \right)^w.
 \end{aligned}$$

Note that in (3.13), the equality holds for $s = 0$ because first, $P_{n,0}^3(w) = 0$ and secondly, the error term $\mathcal{S}_o(w, n, 0)$ (resp. $\mathcal{S}_e(w, n, 0)$) in (3.15) (resp. in (3.17)) is identically zero and therefore, we conclude that $E_{n,0}^3(w) = 0$. \square

Lemma 3.6. *Let γ_1, γ_2 as in Equation (2.10). For $(n, s) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 0}$, and $w \in \mathbb{Z}_{\geq 2}$, then*

$$- \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}} \right)^w \leq - \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n+s}} \right)^w \tag{3.21}$$

and

$$\frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n+s}} \right)^w \leq \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}} \right)^w. \tag{3.22}$$

Proof. The proof of both (3.21) and (3.22) is immediate from the fact that $\frac{1}{\sqrt{n+s}} \leq \frac{1}{\sqrt{n}}$ for all $(n, s) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 0}$. \square

Definition 3.7. Let the coefficient sequence $(g_n)_{n \geq 1}$ be as in Lemma 2.2 and $(\bar{g}_n(s; t))_{n \geq 1}$ be as in Lemma 3.5. Then for $(n, s) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 0}$ and $U \in \mathbb{Z}_{\geq 1}$, we define

$$P_{n,s}(U) := -\log 4\sqrt{3} - \log n + \pi\sqrt{\frac{2n}{3}} + \sum_{u=1}^U \tilde{g}_{u,s} \left(\frac{1}{\sqrt{n}}\right)^u, \tag{3.23}$$

where

$$\tilde{g}_{2u,s} := \frac{(-s)^u}{u} + g_{2u} + \sum_{k=1}^{u-1} \bar{g}_{2k}(s; u) \text{ for all } 1 \leq u \leq \lfloor U/2 \rfloor$$

and

$$\tilde{g}_{2u+1,s} := \pi\sqrt{\frac{2}{3}} \binom{1/2}{u+1} s^{u+1} + g_{2u+1} + \sum_{k=0}^{u-1} \bar{g}_{2k+1}(s; u) \text{ for all } 0 \leq u \leq \lfloor (U-1)/2 \rfloor.$$

Definition 3.8. Let γ_1, γ_2 be as in (2.10). For $(n, s) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 0}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s$, we define

$$E_{n,s}^{\mathcal{U}}(w) := \left(45 \left(s + \frac{1}{24\alpha} \right)^{\lceil \frac{w+1}{2} \rceil} \delta_s + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil}} \right) \frac{1}{w} \left(\frac{1}{\sqrt{n}} \right)^w$$

and

$$E_{n,s}^{\mathcal{L}}(w) := \left(45 \left(s + \frac{1}{24\alpha} \right)^{\lceil \frac{w+1}{2} \rceil} \delta_s + \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil}} \right) \frac{1}{w} \left(\frac{1}{\sqrt{n}} \right)^w.$$

Theorem 3.9. Let $P_{n,s}(U)$ be as in Definition 3.7 and $E_{n,s}^{\mathcal{L}}(w), E_{n,s}^{\mathcal{U}}(w)$ be as in Definition 3.8. If $(n, s) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 0}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > \max\{g(w) - s, 2s\}$, then

$$P_{n,s}(w-1) - E_{n,s}^{\mathcal{L}}(w) < \log p(n+s) < P_{n,s}(w-1) + E_{n,s}^{\mathcal{U}}(w). \tag{3.24}$$

Proof. From (2.9), it follows that for $\lceil \frac{w}{2} \rceil \geq \gamma_0$ and $n > g(w) - s$,

$$\begin{aligned} P_{n+s}(w-1) - \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n+s}} \right)^w &< \log p(n+s) \\ &< P_{n+s}(w-1) + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n+s}} \right)^w, \end{aligned} \tag{3.25}$$

where

$$P_{n+s}(w-1) = -\log 4\sqrt{3} - \log(n+s) + \pi\sqrt{\frac{2(n+s)}{3}} + \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n+s}}\right)^u \quad (\text{by Definition 2.6}).$$

Applying Lemma 3.6 into (3.25), we obtain

$$P_{n+s}(w-1) - \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w < \log p(n+s) < P_{n+s}(w-1) + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w. \tag{3.26}$$

Invoking Lemma 3.3, 3.4, and 3.5 into (3.26), it follows that

$$\begin{aligned} & -\log 4\sqrt{3} + \sum_{i=1}^3 P_{n,s}^i(w) - \sum_{i=1}^3 E_{n,s}^i(w) - \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w < \log p(n+s) \\ & < -\log 4\sqrt{3} + \sum_{i=1}^3 P_{n,s}^i(w) + \sum_{i=1}^3 E_{n,s}^i(w) + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w. \end{aligned} \tag{3.27}$$

For $s \geq 1$,

$$\begin{aligned} & \sum_{i=1}^3 E_{n,s}^i(w) + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w \\ & = \left(\frac{8s^{\lceil \frac{w+1}{2} \rceil}}{\lceil w/2 \rceil} + \frac{29}{w} \left(s + \frac{1}{24\alpha}\right)^{\lceil \frac{w-1}{2} \rceil + 1} + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \right) \left(\frac{1}{\sqrt{n}}\right)^w \\ & = \left(\frac{8s^{\lceil \frac{w+1}{2} \rceil}}{\lceil w/2 \rceil} + \frac{29}{w} \left(s + \frac{1}{24\alpha}\right)^{\lceil \frac{w+1}{2} \rceil} + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \right) \left(\frac{1}{\sqrt{n}}\right)^w \\ & \leq \left(\frac{16s^{\lceil \frac{w+1}{2} \rceil}}{w} + \frac{29}{w} \left(s + \frac{1}{24\alpha}\right)^{\lceil \frac{w+1}{2} \rceil} + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \right) \left(\frac{1}{\sqrt{n}}\right)^w \\ & < \left(45 \left(s + \frac{1}{24\alpha}\right)^{\lceil \frac{w+1}{2} \rceil} + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil}} \right) \frac{1}{w} \left(\frac{1}{\sqrt{n}}\right)^w, \end{aligned} \tag{3.28}$$

and for $s = 0$,

$$\sum_{i=1}^3 E_{n,s}^i(w) + \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w = \frac{\gamma_2}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w. \tag{3.29}$$

Similarly, for $s \geq 1$,

$$\sum_{i=1}^3 E_{n,s}^i(w) + \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w < \left(45 \left(s + \frac{1}{24\alpha}\right)^{\lceil \frac{w+1}{2} \rceil} + \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil}} \right) \frac{1}{w} \left(\frac{1}{\sqrt{n}}\right)^w, \tag{3.30}$$

and for $s = 0$,

$$\sum_{i=1}^3 E_{n,s}^i(w) + \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w = \frac{\gamma_1}{(24\alpha)^{\lceil w/2 \rceil} w} \left(\frac{1}{\sqrt{n}}\right)^w. \tag{3.31}$$

Putting (3.28)-(3.31) together into (3.27), we get

$$-\log 4\sqrt{3} + \sum_{i=1}^3 P_{n,s}^i(w) - E_{n,s}^{\mathcal{L}}(w) < \log p(n+s) < -\log 4\sqrt{3} + \sum_{i=1}^3 P_{n,s}^i(w) + E_{n,s}^{\mathcal{U}}(w). \tag{3.32}$$

From Lemmas 3.3-3.5, it follows that

$$\begin{aligned} & -\log 4\sqrt{3} + \sum_{i=1}^3 P_{n,s}^i(w) \\ &= -\log 4\sqrt{3} - \log n + \pi\sqrt{\frac{2n}{3}} + \sum_{u=1}^{w-1} g_u \left(\frac{1}{\sqrt{n}}\right)^u \\ & \quad + \left(\sum_{k=1}^{\lfloor \frac{w-1}{2} \rfloor} \frac{(-1)^k s^k}{k} \left(\frac{1}{\sqrt{n}}\right)^{2k} + \sum_{t=2}^{\lfloor \frac{w-1}{2} \rfloor} \sum_{u=1}^{t-1} \bar{g}_{2u}(s;t) \left(\frac{1}{\sqrt{n}}\right)^{2t} \right) \\ & \quad + \left(\pi\sqrt{\frac{2}{3}} \sum_{k=1}^{\lfloor \frac{w}{2} \rfloor} \binom{1/2}{k} s^k \left(\frac{1}{\sqrt{n}}\right)^{2k-1} + \sum_{t=1}^{\lfloor \frac{w-2}{2} \rfloor} \sum_{u=0}^{t-1} \bar{g}_{2u+1}(s;t) \left(\frac{1}{\sqrt{n}}\right)^{2t+1} \right) \\ &= -\log 4\sqrt{3} - \log n + \pi\sqrt{\frac{2n}{3}} + \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} \left(\frac{(-s)^u}{u} + g_{2u} + \sum_{k=1}^{u-1} \bar{g}_{2k}(s;u) \right) \left(\frac{1}{\sqrt{n}}\right)^{2u} \\ & \quad + \sum_{u=0}^{\lfloor \frac{w-2}{2} \rfloor} \left(\pi\sqrt{\frac{2}{3}} \binom{1/2}{k+1} s^{k+1} + g_{2u+1} + \sum_{k=0}^{u-1} \bar{g}_{2k+1}(s;u) \right) \left(\frac{1}{\sqrt{n}}\right)^{2u+1} \\ &= -\log 4\sqrt{3} - \log n + \pi\sqrt{\frac{2n}{3}} + \sum_{u=1}^{\lfloor \frac{w-1}{2} \rfloor} \tilde{g}_{2u,s} \left(\frac{1}{\sqrt{n}}\right)^{2u} + \sum_{u=0}^{\lfloor \frac{w-2}{2} \rfloor} \tilde{g}_{2u+1,s} \left(\frac{1}{\sqrt{n}}\right)^{2u+1} \\ &= P_{n,s}(w-1). \end{aligned} \tag{3.33}$$

From (3.32) and (3.33), we conclude the proof of (3.24). \square

Next, we proceed to estimate $\sum_{i=1}^T \log p(n + s_i)$.

Definition 3.10. For $n, T \in \mathbb{Z}_{\geq 1}$ and $\vec{s} := (s_1, s_2, \dots, s_T) \in \mathbb{Z}_{\geq 0}^T$, we define

$$\log p(n; \vec{s}) := \sum_{i=1}^T \log p(n + s_i).$$

Definition 3.11. Let the coefficient sequence $(g_n)_{n \geq 1}$ be as in Lemma 2.2, $(\bar{g}_n(s; t))_{n \geq 1}$ be as in Lemma 3.5, and \vec{s} be as in Definition 3.10. For $n, T \in \mathbb{Z}_{\geq 1}$ and $U \in \mathbb{Z}_{\geq 1}$, we define

$$P_{n, \vec{s}}(U) := -T \cdot \log 4\sqrt{3} - T \cdot \log n + T \cdot \pi \sqrt{\frac{2n}{3}} + \sum_{u=1}^U \tilde{g}_{u, \vec{s}} \left(\frac{1}{\sqrt{n}} \right)^u, \tag{3.34}$$

where

$$\tilde{g}_{2u, \vec{s}} := \frac{1}{u} \sum_{i=1}^T (-s_i)^u + T \cdot g_{2u} + \sum_{i=1}^T \sum_{k=1}^{u-1} \bar{g}_{2k}(s_i; u) \text{ for all } 1 \leq u \leq \lfloor U/2 \rfloor$$

and

$$\tilde{g}_{2u+1, \vec{s}} := \pi \sqrt{\frac{2}{3}} \binom{1/2}{u+1} \sum_{i=1}^T s_i^{u+1} + T \cdot g_{2u+1} + \sum_{i=1}^T \sum_{k=0}^{u-1} \bar{g}_{2k+1}(s_i; u)$$

$$\text{for all } 0 \leq u \leq \lfloor (U-1)/2 \rfloor.$$

Definition 3.12. Let γ_1, γ_2 be as in (2.10) and \vec{s} be as in Definition 3.10. For each $\{s_i\}_{1 \leq i \leq T}$, δ_{s_i} be as in Definition 3.2. For $n, T \in \mathbb{Z}_{\geq 1}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s_i$, we define

$$E_{n, \vec{s}}^{\mathcal{U}}(w) := \left(45 \sum_{i=1}^T \left(s_i + \frac{1}{24\alpha} \right)^{\lceil \frac{w+1}{2} \rceil} \delta_{s_i} + \frac{T \cdot \gamma_2}{(24\alpha)^{\lceil w/2 \rceil}} \right) \frac{1}{w} \left(\frac{1}{\sqrt{n}} \right)^w$$

and

$$E_{n, \vec{s}}^{\mathcal{L}}(w) := \left(45 \sum_{i=1}^T \left(s_i + \frac{1}{24\alpha} \right)^{\lceil \frac{w+1}{2} \rceil} \delta_{s_i} + \frac{T \cdot \gamma_1}{(24\alpha)^{\lceil w/2 \rceil}} \right) \frac{1}{w} \left(\frac{1}{\sqrt{n}} \right)^w.$$

A generalized version of Theorem 3.9 is as follows:

Theorem 3.13. Let $\log p(n; \vec{s})$ be as in Definition 3.10, $P_{n, \vec{s}}(U)$ be as in Definition 3.11, and let $g(k)$ be as in Definition 2.5. Let $E_{n, \vec{s}}^{\mathcal{L}}(w)$ and $E_{n, \vec{s}}^{\mathcal{U}}(w)$ be as in Definition 3.12. If $n, T \in \mathbb{Z}_{\geq 1}$, $w \in \mathbb{Z}_{\geq 2}$, and

$$n > g(w; \vec{s}) := \max_{1 \leq i \leq T} \{g(w) - \min_{1 \leq i \leq T} \{s_i, 2s_i\},$$

then

$$P_{n,\vec{s}}(w - 1) - E_{n,\vec{s}}^{\mathcal{L}}(w) < \log p(n; \vec{s}) < P_{n,\vec{s}}(w - 1) + E_{n,\vec{s}}^{\mathcal{U}}(w). \tag{3.35}$$

Proof. Applying (3.24) for each $\{s_i\}_{1 \leq i \leq T}$ and summing up, we get (3.35). \square

Remark 3.14. A few applications of Theorem 3.13 are listed below.

- (1) Choosing $w = 5$ (resp. $w = 7$), we obtain $(p(n))_{n \geq 26}$ is log-concave (resp. (1.7)).
- (2) Define $u_n := \frac{p(n)p(n+2)}{p(n+1)^2}$ and let N be any positive integer. Then choosing $w = N$, we have a full asymptotic expansion of $\log u_n$ with a precise estimation of the error bound after truncation of the asymptotic expansion at a point N .
- (3) Applying $\vec{s} = \{m, m\}$ and $\vec{r} = \{0, 2m\}$ to (3.35), and estimation of

$$P_{n,\vec{s}}(4) + E_{n,\vec{s}}^{\mathcal{L}}(5) - P_{n,\vec{r}}(4) - E_{n,\vec{s}}^{\mathcal{U}}(5),$$

leads to the strong log-concavity property of $p(n)$.

- (4) Without loss of generality, assume $b = \lambda a$ with $\lambda \geq 1$ in Theorem 1.1. By making the substitutions $(n, \vec{s}) = (a, 0)$, $(n, \vec{s}) = (\lambda a, 0)$, and $(n, \vec{r}) = (a(1 + \lambda), 0)$ to (3.35), we can retrieve (1.10).

4. Asymptotics of $(-1)^{r-1} \Delta^r \log p(n)$

Lemma 4.1. Let $P_{n,s}(w - 1)$ be as in Theorem 3.9. Then for all $r \geq 2$,

$$\sum_{i=0}^r \binom{r}{i} (-1)^{i+1} P_{n,i}(2r) = C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1} - (r - 1)! \left(\frac{1}{\sqrt{n}}\right)^{2r}, \tag{4.1}$$

where $C_r = \frac{\pi}{\sqrt{6}} \left(\frac{1}{2}\right)_{r-1}$ and $(a)_k$ is the standard notation for the rising factorial.

Proof. From Definition 3.7, it follows that

$$\begin{aligned} & \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} P_{n,i}(2r) \\ &= \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \left(-\log 4\sqrt{3} - \log n + \sqrt{\frac{2n}{3}} + \sum_{u=1}^{2r} \tilde{g}_{u,i} \left(\frac{1}{\sqrt{n}}\right)^u \right) \\ &= \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \sum_{u=1}^{2r} \tilde{g}_{u,i} \left(\frac{1}{\sqrt{n}}\right)^u \\ &= \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \sum_{u=1}^{2r-2} \tilde{g}_{u,i} \left(\frac{1}{\sqrt{n}}\right)^u + \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \tilde{g}_{2r-1,i} \left(\frac{1}{\sqrt{n}}\right)^{2r-1} \end{aligned}$$

$$+ \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \tilde{g}_{2r,i} \left(\frac{1}{\sqrt{n}}\right)^{2r}. \tag{4.2}$$

Following the notation from [11], here $\left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\}$ denotes the Stirling number of second kind which counts the number of ways to partition a set of n elements into m nonempty subsets. Here we state three facts about $\left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\}$ which is required in the later part of the proof:

Fact 4.2. [11, Table 264] $\left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\} = \binom{n}{m} = 0$.

Fact 4.3. [11, Table 264] $\left\{ \begin{smallmatrix} n \\ n \end{smallmatrix} \right\} = \binom{n}{n} = 1$.

Fact 4.4. [11, Table 265, Eq. (6.19)] $m! \left\{ \begin{smallmatrix} n \\ m \end{smallmatrix} \right\} = \sum_{k=0}^m \binom{m}{k} k^n (-1)^{m-k}$.

For all integers $1 \leq u \leq 2r - 2$ and $u \equiv 0 \pmod{2}$, we have

$$\begin{aligned} & \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \sum_{u=1}^{r-1} \tilde{g}_{2u,i} \left(\frac{1}{\sqrt{n}}\right)^{2u} \\ &= \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \sum_{u=1}^{r-1} \left[\frac{(-i)^u}{u} + g_{2u} + \sum_{k=1}^{u-1} \bar{g}_{2k}(i; u) \right] \left(\frac{1}{\sqrt{n}}\right)^{2u} \\ &= \sum_{u=1}^{r-1} \frac{(-1)^u}{u} (-1)^{r+1} r! \left\{ \begin{smallmatrix} u \\ r \end{smallmatrix} \right\} \left(\frac{1}{\sqrt{n}}\right)^{2u} \quad (\text{by Fact 4.4}) \\ & \quad + \sum_{u=1}^{r-1} \sum_{k=1}^{u-1} g_{2k} \binom{-k}{u-k} \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} i^{u-k} \left(\frac{1}{\sqrt{n}}\right)^{2u} \\ &= \sum_{u=1}^{r-1} \frac{(-1)^u}{u} (-1)^{r+1} r! \left\{ \begin{smallmatrix} u \\ r \end{smallmatrix} \right\} \left(\frac{1}{\sqrt{n}}\right)^{2u} \\ & \quad + \sum_{u=1}^{r-1} \sum_{k=1}^{u-1} g_{2k} \binom{-k}{u-k} (-1)^{r+1} r! \left\{ \begin{smallmatrix} u-k \\ r \end{smallmatrix} \right\} \left(\frac{1}{\sqrt{n}}\right)^{2u} \\ &= 0 \quad (\text{by Fact 4.2}). \end{aligned} \tag{4.3}$$

Similarly for all integers $1 \leq u \leq 2r - 2$ and $u \equiv 1 \pmod{2}$, we obtain

$$\begin{aligned} & \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \sum_{u=0}^{r-2} \tilde{g}_{2u+1,i} \left(\frac{1}{\sqrt{n}}\right)^{2u+1} \\ &= \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \sum_{u=0}^{r-2} \left[\pi \sqrt{\frac{2}{3}} \binom{1/2}{u+1} i^{u+1} + g_{2u+1} + \sum_{k=0}^{u-1} \bar{g}_{2k+1}(i; u) \right] \left(\frac{1}{\sqrt{n}}\right)^{2u+1} \end{aligned}$$

$$\begin{aligned}
 &= \sum_{u=0}^{r-2} \pi \sqrt{\frac{2}{3}} \binom{1/2}{u+1} (-1)^{r+1} r! \left\{ \begin{matrix} u+1 \\ r \end{matrix} \right\} \left(\frac{1}{\sqrt{n}} \right)^{2u+1} \\
 &+ \sum_{u=0}^{r-2} \sum_{k=0}^{u-1} g_{2k+1} \binom{-k-1/2}{u-k} (-1)^{r+1} r! \left\{ \begin{matrix} u-k \\ r \end{matrix} \right\} \left(\frac{1}{\sqrt{n}} \right)^{2u+1} \quad (\text{by Fact 4.4}) \\
 &= 0 \quad (\text{by Fact 4.2}).
 \end{aligned} \tag{4.4}$$

From (4.3) and (4.4), it follows that for all $1 \leq u \leq 2r - 2$,

$$\sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \sum_{u=1}^{2r-2} \tilde{g}_{u,i} \left(\frac{1}{\sqrt{n}} \right)^u = 0.$$

Now,

$$\begin{aligned}
 &\sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \tilde{g}_{2r-1,i} \left(\frac{1}{\sqrt{n}} \right)^{2r-1} \\
 &= \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \left[\pi \sqrt{\frac{2}{3}} \binom{1/2}{r} i^r + g_{2r-1} + \sum_{k=0}^{r-2} \bar{g}_{2k+1}(i; r-1) \right] \left(\frac{1}{\sqrt{n}} \right)^{2r-1} \\
 &= \left[\pi \sqrt{\frac{2}{3}} \binom{1/2}{r} (-1)^{r+1} r! \left\{ \begin{matrix} r \\ r \end{matrix} \right\} \right. \\
 &\quad \left. + \sum_{k=0}^{r-2} g_{2k+1} \binom{-k-1/2}{r-1-k} (-1)^{r+1} r! \left\{ \begin{matrix} r-1-k \\ r \end{matrix} \right\} \right] \left(\frac{1}{\sqrt{n}} \right)^{2r-1} \\
 &= \frac{\pi}{\sqrt{6}} \left(\frac{1}{2} \right)_{r-1} \left(\frac{1}{\sqrt{n}} \right)^{2r-1} \quad (\text{by Facts 4.2 - 4.4}).
 \end{aligned} \tag{4.5}$$

We finish the proof by showing that

$$\begin{aligned}
 &\sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \tilde{g}_{2r,i} \left(\frac{1}{\sqrt{n}} \right)^{2r} \\
 &= \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \left[\frac{(-i)^r}{r} + g_{2r} + \sum_{k=1}^{r-1} \bar{g}_{2k}(i; r) \right] \left(\frac{1}{\sqrt{n}} \right)^{2r} \\
 &= \left[-(r-1)! + \sum_{k=1}^{r-1} g_{2k} \binom{-k}{r-k} (-1)^{r+1} r! \left\{ \begin{matrix} r-k \\ r \end{matrix} \right\} \right] \left(\frac{1}{\sqrt{n}} \right)^{2r} \\
 &= -(r-1)! \left(\frac{1}{\sqrt{n}} \right)^{2r-1}. \quad \square
 \end{aligned} \tag{4.6}$$

Definition 4.5. Let γ_1 be as in (2.10) and C_r be as in Lemma 4.1. Then for all $r \geq 2$, define

$$L_1(r) := \left(\frac{\gamma_1}{(12\alpha)^{r+1}} + 45 \sum_{i=1}^r \binom{r}{i} \left(i + \frac{1}{24\alpha} \right)^{r+1} \right) \frac{1}{2r+1},$$

$$L(r) := (r-1)! + L_1(r),$$

and

$$N_L(r) := \max \left\{ \left(\frac{L(r)}{C_r} \right)^2, g(2r+1) \right\}.$$

Lemma 4.6. *Let $L(r), N_L(r)$ be as in Definition 4.5 and C_r be as in Lemma 4.1. Then for all $n > N_L(r)$,*

$$(-1)^{r-1} \Delta^r \log p(n) > \log \left(1 + C_r \left(\frac{1}{\sqrt{n}} \right)^{2r-1} - L(r) \left(\frac{1}{\sqrt{n}} \right)^{2r} \right). \tag{4.7}$$

Proof. We split $(-1)^{r-1} \Delta^r \log p(n)$ as follows:

$$\begin{aligned} (-1)^{r-1} \Delta^r \log p(n) &= \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \log p(n+i) \\ &= \sum_{i=0}^{\lfloor \frac{r-1}{2} \rfloor} \binom{r}{2i+1} \log p(n+2i+1) - \sum_{i=0}^{\lfloor \frac{r}{2} \rfloor} \binom{r}{2i} \log p(n+2i). \end{aligned} \tag{4.8}$$

Applying Theorem 3.9 with $w = 2r + 1$ to (4.8), for all $n > g(2r + 1) = \max_{0 \leq i \leq r} \{g(2r + 1) - i, 2i\}$, we have

$$\begin{aligned} &(-1)^{r-1} \Delta^r \log p(n) \\ &> \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} P_{n,i}(2r) - \sum_{i=0}^{\lfloor \frac{r-1}{2} \rfloor} \binom{r}{2i+1} E_{n,2i+1}^{\mathcal{L}}(2r+1) - \sum_{i=0}^{\lfloor \frac{r}{2} \rfloor} \binom{r}{2i} E_{n,2i}^{\mathcal{U}}(2r+1) \\ &= C_r \left(\frac{1}{\sqrt{n}} \right)^{2r-1} - (r-1)! \left(\frac{1}{\sqrt{n}} \right)^{2r} - \sum_{i=0}^{\lfloor \frac{r-1}{2} \rfloor} \binom{r}{2i+1} E_{n,2i+1}^{\mathcal{L}}(2r+1) \\ &\quad - \sum_{i=0}^{\lfloor \frac{r}{2} \rfloor} \binom{r}{2i} E_{n,2i}^{\mathcal{U}}(2r+1) \quad (\text{by Lemma 4.1}). \end{aligned} \tag{4.9}$$

From Definition 3.8, it is clear that $E_{n,s}^{\mathcal{U}}(w) < E_{n,s}^{\mathcal{L}}(w)$ because $\gamma_2 < \gamma_1$. Therefore,

$$\sum_{i=0}^{\lfloor \frac{r-1}{2} \rfloor} \binom{r}{2i+1} E_{n,2i+1}^{\mathcal{L}}(2r+1) + \sum_{i=0}^{\lfloor \frac{r}{2} \rfloor} \binom{r}{2i} E_{n,2i}^{\mathcal{U}}(2r+1) < \sum_{i=0}^r \binom{r}{i} E_{n,i}^{\mathcal{L}}(2r+1), \tag{4.10}$$

and

$$\sum_{i=0}^r \binom{r}{i} E_{n,i}^{\mathcal{L}}(2r+1) = L_1(r) \left(\frac{1}{\sqrt{n}}\right)^{2r+1}. \tag{4.11}$$

From (4.9) and (4.11), it follows that

$$\begin{aligned} (-1)^{r-1} \Delta^r \log p(n) &> C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1} - (r-1)! \left(\frac{1}{\sqrt{n}}\right)^{2r} - L_1(r) \left(\frac{1}{\sqrt{n}}\right)^{2r+1} \\ &> C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1} - L(r) \left(\frac{1}{\sqrt{n}}\right)^{2r}, \end{aligned} \tag{4.12}$$

and consequently for all $n > N_L(r)$, we get

$$(-1)^{r-1} \Delta^r \log p(n) > \log \left(1 + C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1} - L(r) \left(\frac{1}{\sqrt{n}}\right)^{2r} \right). \quad \square$$

Definition 4.7. Let $L_1(r)$ be as in Definition 4.5 and C_r be as in Lemma 4.1. Then for all $r \geq 2$, define

$$N_U(r) := \max \left\{ \left(\frac{L_1(r)+1}{(r-1)!} \right)^2, \left(\frac{C_r^2}{2} \right)^{2/2r-3}, g(2r+1) \right\}.$$

Lemma 4.8. Let $L_1(r)$ be as in Definition 4.5, C_r be as in Lemma 4.1, and $N_U(r)$ be as in Definition 4.7. Then for all $n > N_U(r)$,

$$(-1)^{r-1} \Delta^r \log p(n) < \log \left(1 + C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1} \right). \tag{4.13}$$

Proof. Applying Theorem 3.9 with $w = 2r + 1$ to (4.8), we have for all $n > g(2r + 1)$,

$$\begin{aligned} &(-1)^{r-1} \Delta^r \log p(n) \\ &< \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} P_{n,i}(2r) + \sum_{i=0}^{\lfloor \frac{r-1}{2} \rfloor} \binom{r}{2i+1} E_{n,2i+1}^{\mathcal{U}}(2r+1) + \sum_{i=0}^{\lfloor \frac{r}{2} \rfloor} \binom{r}{2i} E_{n,2i}^{\mathcal{L}}(2r+1) \\ &< C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1} - (r-1)! \left(\frac{1}{\sqrt{n}}\right)^{2r} + \sum_{i=0}^r \binom{r}{i} E_{n,i}^{\mathcal{L}}(2r+1) \quad (\text{by Lemma 4.1}) \\ &= C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1} - (r-1)! \left(\frac{1}{\sqrt{n}}\right)^{2r} + L_1(r) \left(\frac{1}{\sqrt{n}}\right)^{2r+1}. \end{aligned} \tag{4.14}$$

For all $n > N_U(r)$, it follows that

$$-(r-1)! \left(\frac{1}{\sqrt{n}}\right)^{2r} + L_1(r) \left(\frac{1}{\sqrt{n}}\right)^{2r+1} < -\frac{C_r^2}{2n^{2r-1}}. \tag{4.15}$$

From (4.14) and (4.15), it follows that for all $n > N_U(r)$,

$$(-1)^{r-1} \Delta^r \log p(n) < \log \left(1 + C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1}\right). \quad \square$$

Theorem 4.9. *Let $L(r), N_L(r)$ be as in Definition 4.5 and $N_U(r)$ be as in Definition 4.7. Let C_r be as in Lemma 4.1. Then for all $n > N(r) := \max\{N_L(r), N_U(r)\}$,*

$$\log \left(1 + C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1} - L(r) \left(\frac{1}{\sqrt{n}}\right)^{2r}\right) < (-1)^{r-1} \Delta^r \log p(n) < \log \left(1 + C_r \left(\frac{1}{\sqrt{n}}\right)^{2r-1}\right). \tag{4.16}$$

Proof. Lemmas 4.1 and 4.6 together imply (4.16). \square

Theorem 4.10. *For all $r \geq 2$,*

$$(-1)^{r-1} \Delta^r \log p(n) \underset{n \rightarrow \infty}{\sim} \sum_{u=2r-1}^{\infty} G_u \left(\frac{1}{\sqrt{n}}\right)^u, \tag{4.17}$$

with

$$\begin{aligned} G_{2u} &= \left[\frac{(-1)^u}{u} \begin{Bmatrix} u \\ r \end{Bmatrix} + \sum_{k=1}^{u-r} g_{2k} \begin{Bmatrix} -k \\ u-k \end{Bmatrix} \begin{Bmatrix} u-k \\ r \end{Bmatrix} \right] (-1)^{r+1} r! \text{ for all } u \geq r, \\ G_{2u+1} &= \left[\pi \sqrt{\frac{2}{3}} \begin{Bmatrix} 1/2 \\ u+1 \end{Bmatrix} \begin{Bmatrix} u+1 \\ r \end{Bmatrix} + \sum_{k=0}^{u-r} g_{2k+1} \begin{Bmatrix} -k-1/2 \\ u-k \end{Bmatrix} \begin{Bmatrix} u-k \\ r \end{Bmatrix} \right] (-1)^{r+1} r! \\ &\text{for all } u \geq r-1. \end{aligned} \tag{4.18}$$

Proof. Following (4.2) and letting $w \rightarrow \infty$, we obtain

$$(-1)^{r-1} \Delta^r \log p(n) \underset{n \rightarrow \infty}{\sim} \sum_{u=2r-1}^{\infty} \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \tilde{g}_{u,i} \left(\frac{1}{\sqrt{n}}\right)^u. \tag{4.19}$$

For all $u \geq 2r-1$ and $u \equiv 0 \pmod{2}$, we get

$$\begin{aligned} \sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \tilde{g}_{2u,i} &= \left[\frac{(-1)^u}{u} \begin{Bmatrix} u \\ r \end{Bmatrix} + \sum_{k=1}^{u-1} g_{2k} \begin{Bmatrix} -k \\ u-k \end{Bmatrix} \begin{Bmatrix} u-k \\ r \end{Bmatrix} \right] (-1)^{r+1} r! \\ &= \left[\frac{(-1)^u}{u} \begin{Bmatrix} u \\ r \end{Bmatrix} + \sum_{k=1}^{u-r} g_{2k} \begin{Bmatrix} -k \\ u-k \end{Bmatrix} \begin{Bmatrix} u-k \\ r \end{Bmatrix} \right] (-1)^{r+1} r!. \end{aligned}$$

Similarly, for all $u \geq 2r - 1$ and $u \equiv 1 \pmod{2}$, it follows that

$$\sum_{i=0}^r \binom{r}{i} (-1)^{i+1} \tilde{g}_{2u+1,i} = \left[\pi \sqrt{\frac{2}{3}} \binom{1/2}{u+1} \left\{ \begin{matrix} u+1 \\ r \end{matrix} \right\} + \sum_{k=0}^{u-r} g_{2k+1} \right. \\ \left. \times \binom{-k-1/2}{u-k} \left\{ \begin{matrix} u-k \\ r \end{matrix} \right\} \right] (-1)^{r+1} r!. \quad \square$$

5. A framework to verify multiplicative inequalities for $p(n)$

Here we list down the steps in order to make a decision whether a given multiplicative inequality holds or not.

- (Step 0): Given $\prod_{i=1}^T p(n + s_i)$ and $\prod_{i=1}^T p(n + r_i)$ with $T \geq 1$. Without loss of generality, assume that s_i, r_i are non-negative integers for all $1 \leq i \leq T$. Transform the products into additive ones by applying the natural logarithm; i.e., $\sum_{i=1}^T \log p(n + s_i)$ and $\sum_{i=1}^T \log p(n + r_i)$.
- (Step 1): Choose $w = m + 1$, where $(s_1, \dots, s_T) \stackrel{m}{=} (r_1, \dots, r_T)$. From (3.35), we observe that for each $1 \leq i \leq T$, $\log p(n + s_i)$ and $\log p(n + r_i)$ has the main term $P_{n,\bar{s}}(w - 1)$ and $P_{n,\bar{r}}(w - 1)$ respectively. Consequently, each of these main terms are dominated by $T \cdot c \sum_{i=1}^T \sqrt{n + s_i}$ and $T \cdot c \sum_{i=1}^T \sqrt{n + r_i}$ with $c = \pi \sqrt{2/3}$ respectively. Therefore, in order to choose w , it is enough to compute the Taylor expansion of $\sum_{i=1}^T (\sqrt{n + s_i} - \sqrt{n + r_i})$ which is given by:

$$\sum_{i=1}^T (\sqrt{n + s_i} - \sqrt{n + r_i}) = \sum_{m=1}^{\infty} \frac{\binom{1/2}{m}}{\sqrt{n}^{2m-1}} \sum_{i=1}^T (s_i^m - r_i^m). \tag{5.1}$$

So our optimal choice is such minimal $m \geq 1$ so that $\sum_{i=1}^T (s_i^m - r_i^m) \neq 0$.

- (Step 2): Applying $w = m + 1$ as in the previous step to Theorem (3.35), it remains to verify whether

$$P_{n,\bar{s}}(m) - E_{n,\bar{s}}^{\mathcal{L}}(m + 1) > P_{n,\bar{r}}(m) + E_{n,\bar{r}}^{\mathcal{U}}(m + 1) \tag{5.2}$$

or

$$P_{n,\vec{r}}(m) - E_{n,\vec{r}}^{\mathcal{L}}(m+1) > P_{n,\vec{s}}(m) + E_{n,\vec{s}}^{\mathcal{U}}(m+1), \tag{5.3}$$

in order to decide whether $\sum_{i=1}^T \log p(n + s_i) \geq \sum_{i=1}^T \log p(n + r_i)$ or $\sum_{i=1}^T \log p(n + r_i) \geq \sum_{i=1}^T \log p(n + s_i)$ respectively.

6. Inequalities for $p(n; \vec{s})$

Definition 6.1. Let $\tilde{g}_{u,\vec{s}}$ be as in Definition 3.11, and \vec{s} be as in Definition 3.10. For $n, T, U \in \mathbb{Z}_{\geq 1}$, define

$$\mathcal{M}(n; T) := \left(\frac{e^{\pi\sqrt{2n/3}}}{4n\sqrt{3}} \right)^T,$$

and

$$\tilde{P}_{n,\vec{s}}(U) := \exp\left(\sum_{u=1}^U \tilde{g}_{u,\vec{s}}\left(\frac{1}{\sqrt{n}}\right)^u\right).$$

Definition 6.2. Let γ_1, γ_2 be as in (2.10) and \vec{s} be as in Definition 3.10. For each $\{s_i\}_{1 \leq i \leq T}$, δ_{s_i} be as in Definition 3.2. For $n, T \in \mathbb{Z}_{\geq 1}$, $w \in \mathbb{Z}_{\geq 2}$, and $n > 2s_i$, we define

$$C_{\mathcal{U}}(w; \vec{s}) := \left(45 \sum_{i=1}^T \left(s_i + \frac{1}{24\alpha}\right)^{\lceil \frac{w+1}{2} \rceil} \delta_{s_i} + \frac{T \cdot \gamma_2}{(24\alpha)^{\lceil w/2 \rceil}} \right) \frac{1}{w}$$

and

$$C_{\mathcal{L}}(w; \vec{s}) := \left(45 \sum_{i=1}^T \left(s_i + \frac{1}{24\alpha}\right)^{\lceil \frac{w+1}{2} \rceil} \delta_{s_i} + \frac{T \cdot \gamma_1}{(24\alpha)^{\lceil w/2 \rceil}} \right) \frac{1}{w}.$$

Lemma 6.3. Let $\log p(n; \vec{s})$ be as in Definition 3.10, and let $g(k)$ be as in Definition 2.5. Let $\mathcal{M}(n; T)$ and $\tilde{P}_{n,\vec{s}}(U)$ be as in Definition 6.1. Let $g(w; \vec{s})$ be as in Theorem 3.13, and $C_{\mathcal{L}}(w; \vec{s}), C_{\mathcal{U}}(w; \vec{s})$ be as in Definition 6.2. If $n, T \in \mathbb{Z}_{\geq 1}$, $w \in \mathbb{Z}_{\geq 2}$, and

$$n > N_1(w; \vec{s}) := \max\left\{g(w; \vec{s}), \left(C_{\mathcal{L}}(w; \vec{s})\right)^{2/w}, \left(C_{\mathcal{U}}(w; \vec{s})\right)^{2/w}\right\},$$

then

$$\mathcal{M}(n; T) \tilde{P}_{n,\vec{s}}(w-1) \left(1 - C_{\mathcal{L}}(w; \vec{s}) \left(\frac{1}{\sqrt{n}}\right)^w\right)$$

$$< p(n; \vec{s}) < \mathcal{M}(n; T) \tilde{P}_{n, \vec{s}}(w-1) \left(1 + 2 C_U(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w \right). \tag{6.1}$$

Proof. Applying the exponential function on both sides of the inequality (3.35), we get for all $n > g(w; \vec{s})$,

$$\mathcal{M}(n; T) \tilde{P}_{n, \vec{s}}(w-1) e^{-E_{n, \vec{s}}^{\mathcal{L}}(w)} < p(n; \vec{s}) < \mathcal{M}(n; T) \tilde{P}_{n, \vec{s}}(w-1) e^{E_{n, \vec{s}}^{\mathcal{U}}(w)}. \tag{6.2}$$

Now for all $n > \max \left\{ \left(C_{\mathcal{L}}(w; \vec{s}) \right)^{2/w}, C_U(w; \vec{s}) \right\}^{2/w}$, it follows that

$$0 < E_{n, \vec{s}}^{\mathcal{U}}(w) < 1 \text{ and } 0 < E_{n, \vec{s}}^{\mathcal{L}}(w) < 1. \tag{6.3}$$

For all $0 < x < 1$, we know that $e^x < 1 + 2x$ and $e^{-x} > 1 - x$. Therefore from (6.3) and following Definition 3.12, we finally have

$$e^{E_{n, \vec{s}}^{\mathcal{U}}(w)} < 1 + 2 C_U(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w \text{ and } e^{-E_{n, \vec{s}}^{\mathcal{L}}(w)} > 1 - C_{\mathcal{L}}(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w. \tag{6.4}$$

Equations (6.2) and (6.4) together imply (6.1). \square

Definition 6.4. For $k \in \mathbb{Z}_{\geq 0}$, $w \geq 2$, and $\vec{\ell} := (\ell_1, \dots, \ell_{w-1})$, define

$$\begin{aligned}
 X(k) &:= \left\{ \vec{\ell} \in \mathbb{Z}_{\geq 0}^{w-1} : \sum_{u=1}^{w-1} \ell_u = k \right\}, \\
 X_{\mathcal{M}}(k) &:= \left\{ \vec{\ell} \in X(k) : 0 \leq \sum_{u=1}^{w-1} u \ell_u \leq w-1 \right\},
 \end{aligned}$$

and

$$X_{\mathcal{E}}(k) := \left\{ \vec{\ell} \in X(k) : \sum_{u=1}^{w-1} u \ell_u \geq w \right\}.$$

Definition 6.5. Let $X(k)$ and $X_{\mathcal{M}}(k)$ be as in Definition 6.4 and $\tilde{g}_{u, \vec{s}}$ be as in Definition 3.11. Then for all $w \geq 2$, define

$$\hat{P}_{n, \vec{s}}(w-1) := \sum_{k=0}^{w-1} \frac{1}{k!} \sum_{\vec{\ell} \in X_{\mathcal{M}}(k)} F(k; w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^{\sum_{u=1}^{w-1} u \ell_u},$$

and

$$\widehat{E}_{n,\vec{s}}(w-1) := \sum_{k=0}^{w-1} \frac{1}{k!} \sum_{\vec{\ell} \in X_{\mathcal{E}}(k)} F(k; w; \vec{s}) \left(\frac{1}{\sqrt{n}}\right)^{\sum_{u=1}^{w-1} u\ell_u},$$

where

$$F(k; w; \vec{s}) := \binom{k}{\ell_1, \dots, \ell_{w-1}} \prod_{u=1}^{w-1} (\tilde{g}_{u,\vec{s}})^{\ell_u},$$

with $\binom{k}{\ell_1, \dots, \ell_{w-1}} = \frac{k!}{\ell_1! \dots \ell_{w-1}!}$ is a multinomial coefficient.

Definition 6.6. Let $X_{\mathcal{E}}(k)$ be as in Definition 6.4 and $F(k; w; \vec{s})$ be as in Definition 6.5 and $\tilde{g}_{u,\vec{s}}$ be as in Definition 3.11. For $w \geq 2$, define

$$E(w; \vec{s}) := \sum_{k=0}^{w-1} \frac{1}{k!} \sum_{\vec{\ell} \in X_{\mathcal{E}}(k)} \left| F(k; w; \vec{s}) \right| + 3 \left(|\tilde{g}_{1,\vec{s}}| + 1 \right)^w.$$

Lemma 6.7. Let $\tilde{P}_{n,\vec{s}}(U)$ be as in Definition 6.1 and $X_{\mathcal{E}}(k)$ be as in Definition 6.4. Let $\widehat{P}_{n,\vec{s}}(w-1)$, $\widehat{P}_{n,\vec{s}}(w-1)$, and $F(k; w; \vec{s})$ be as in Definition 6.5. Let $E(w; \vec{s})$ be as in Definition 6.6. Then for all $w \geq 2$ and

$$n > N_2(w; \vec{s}) := \max_{1 \leq u \leq w-1} \left\{ \left((w-1) |\tilde{g}_{u,\vec{s}}| \right)^{2/u} \right\},$$

we have

$$\left| \tilde{P}_{n,\vec{s}}(w-1) - \widehat{P}_{n,\vec{s}}(w-1) \right| < E(w; \vec{s}) \left(\frac{1}{\sqrt{n}}\right)^w. \tag{6.5}$$

Proof. Expanding $\tilde{P}_{n,\vec{s}}(w-1)$ and splitting it as follows:

$$\begin{aligned} \tilde{P}_{n,\vec{s}}(w-1) &= \widehat{P}_{n,\vec{s}}(w-1) + \widehat{E}_{n,\vec{s}}(w-1) + \sum_{k=w}^{\infty} \frac{1}{k!} \sum_{\vec{\ell} \in X(k)} F(k; w; \vec{s}) \left(\frac{1}{\sqrt{n}}\right)^{\sum_{u=1}^{w-1} u\ell_u} \\ &= \widehat{P}_{n,\vec{s}}(w-1) + \widehat{E}_{n,\vec{s}}(w-1) + \sum_{k=w}^{\infty} \frac{1}{k!} \left(\sum_{u=1}^{w-1} \frac{\tilde{g}_{u,\vec{s}}}{\sqrt{n}^u} \right)^k. \end{aligned} \tag{6.6}$$

Therefore,

$$\left| \tilde{P}_{n,\vec{s}}(w-1) - \widehat{P}_{n,\vec{s}}(w-1) \right|$$

$$\begin{aligned}
 &\leq \left| \widehat{E}_{n,\vec{s}}(w-1) \right| + \left(\sum_{u=1}^{w-1} \frac{|\tilde{g}_{u,\vec{s}}|}{\sqrt{n}^u} \right)^w \sum_{k=0}^{\infty} \frac{1}{(k+w)!} \left(\sum_{u=1}^{w-1} \frac{|\tilde{g}_{u,\vec{s}}|}{\sqrt{n}^u} \right)^k \\
 &= \left| \widehat{E}_{n,\vec{s}}(w-1) \right| + \left(\frac{1}{\sqrt{n}} \right)^w \left(|\tilde{g}_{1,\vec{s}}| + \sum_{u=1}^{w-2} \frac{|\tilde{g}_{u+1,\vec{s}}|}{\sqrt{n}^u} \right)^w \sum_{k=0}^{\infty} \frac{1}{(k+w)!} \left(\sum_{u=1}^{w-1} \frac{|\tilde{g}_{u,\vec{s}}|}{\sqrt{n}^u} \right)^k \\
 &< \left| \widehat{E}_{n,\vec{s}}(w-1) \right| + \left(\frac{1}{\sqrt{n}} \right)^w (|\tilde{g}_{1,\vec{s}}| + 1)^w \sum_{k=0}^{\infty} \frac{1}{(k+w)!} \quad (\text{since } n > N_2(w; \vec{s})) \\
 &\leq \left| \widehat{E}_{n,\vec{s}}(w-1) \right| + \frac{(|\tilde{g}_{1,\vec{s}}| + 1)^w}{w!} \left(\frac{1}{\sqrt{n}} \right)^w \sum_{k=0}^{\infty} \frac{1}{k!} \\
 &< \left| \widehat{E}_{n,\vec{s}}(w-1) \right| + 3 \frac{(|\tilde{g}_{1,\vec{s}}| + 1)^w}{w!} \left(\frac{1}{\sqrt{n}} \right)^w. \tag{6.7}
 \end{aligned}$$

Now,

$$\begin{aligned}
 \left| \widehat{E}_{n,\vec{s}}(w-1) \right| &\leq \sum_{k=0}^{w-1} \frac{1}{k!} \sum_{\vec{\ell} \in X_{\mathcal{E}}(k)} \left| F(k; w; \vec{s}) \right| \left(\frac{1}{\sqrt{n}} \right)^{\sum_{u=1}^{w-1} u \ell_u} \\
 &\leq \sum_{k=0}^{w-1} \frac{1}{k!} \sum_{\vec{\ell} \in X_{\mathcal{E}}(k)} \left| F(k; w; \vec{s}) \right| \left(\frac{1}{\sqrt{n}} \right)^w \quad (\text{since } \vec{\ell} \in X_{\mathcal{E}}(k)). \tag{6.8}
 \end{aligned}$$

Combining (6.7) and (6.8), we get (6.5). \square

Definition 6.8. Let $C_{\mathcal{U}}(w; \vec{s})$ and $C_{\mathcal{L}}(w; \vec{s})$ be as in Definition 6.2. Let $E(w; \vec{s})$ be as in Definition 6.6. Then for all $w \geq 2$, define

$$E_L(w; \vec{s}) := 3 C_{\mathcal{L}}(w; \vec{s}) + E(w; \vec{s}),$$

and

$$E_U(w; \vec{s}) := 6 C_{\mathcal{U}}(w; \vec{s}) + E(w; \vec{s}) \left(2 C_{\mathcal{U}}(w; \vec{s}) + 1 \right).$$

Theorem 6.9. Let $\mathcal{M}(n; T)$ be as in Definition 6.1 and $\widehat{P}_{n,\vec{s}}(w-1)$ be as in Definition 6.5. Let $E_{n,\vec{s}}^L(w)$ and $E_{n,\vec{s}}^U(w)$ be as in Definition 6.8. Let $N_1(w; \vec{s})$ and $N_2(w; \vec{s})$ be as in Lemmas 6.3 and 6.7. Then for all $w \geq 2$ and

$$N > N(w; \vec{s}) := \max \left\{ N_1(w; \vec{s}), N_2(w; \vec{s}) \right\},$$

we have

$$\begin{aligned} & \mathcal{M}(n; T) \left(\widehat{P}_{n, \vec{s}}(w-1) - E_L(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w \right) \\ & < p(n; \vec{s}) < \mathcal{M}(n; T) \left(\widehat{P}_{n, \vec{s}}(w-1) + E_U(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w \right). \end{aligned} \tag{6.9}$$

Proof. From Lemmas 6.3 and 6.7, for $n > N(w; \vec{s})$, it follows that

$$p(n; \vec{s}) < \mathcal{M}(n; T) \left(\widehat{P}_{n, \vec{s}}(w-1) + E(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w \right) \left(1 + 2 C_U(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w \right), \tag{6.10}$$

and

$$p(n; \vec{s}) > \mathcal{M}(n; T) \left(\widehat{P}_{n, \vec{s}}(w-1) - E(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w \right) \left(1 - C_L(w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^w \right). \tag{6.11}$$

Now

$$\begin{aligned} \left| \widehat{P}_{n, \vec{s}}(w-1) \right| &= \left| \sum_{k=0}^{w-1} \frac{1}{k!} \sum_{\vec{\ell} \in X_{\mathcal{M}}(k)} F(k; w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^{\sum_{u=1}^{w-1} u \ell_u} \right| \\ &\leq \left| \sum_{k=0}^{w-1} \frac{1}{k!} \sum_{\vec{\ell} \in X(k)} F(k; w; \vec{s}) \left(\frac{1}{\sqrt{n}} \right)^{\sum_{u=1}^{w-1} u \ell_u} \right| \quad \left(\text{as } X_{\mathcal{M}}(k) \subseteq X(k) \right) \\ &= \left| \sum_{k=0}^{w-1} \frac{1}{k!} \left(\sum_{u=1}^{w-1} \frac{\tilde{g}_{u, \vec{s}}}{\sqrt{n}^u} \right)^k \right| \leq \sum_{k=0}^{w-1} \frac{1}{k!} \left(\sum_{u=1}^{w-1} \frac{|\tilde{g}_{u, \vec{s}}|}{\sqrt{n}^u} \right)^k \\ &< \sum_{k=0}^{w-1} \frac{1}{k!} \quad \left(\text{as } n > N_2(w; \vec{s}) \right) < 3. \end{aligned} \tag{6.12}$$

Applying (6.12) to (6.10), we arrive at the upper bound of (6.9). We get the lower bound of (6.9) by applying (6.12) to (6.11) and from the fact that $C_L(w; \vec{s}) \cdot E(w; \vec{s}) > 0$ for all $w \geq 2$. \square

7. Conclusion

We conclude this paper by pointing out the following aspects in which Theorem 6.9 remains incomplete.

- (1) Suppose we are given the following two functions defined by shifts of $p(n)$:

$$SP(n; S) := \sum_{j=1}^M \prod_{i=1}^T p(n + s_{i,j}) \quad \text{and} \quad SP(n; R) := \sum_{j=1}^M \prod_{i=1}^T p(n + r_{i,j}),$$

where $S = (s_{i,j})_{1 \leq i \leq T, 1 \leq j \leq M}$ and $R = (r_{i,j})_{1 \leq i \leq T, 1 \leq j \leq M}$. Now in order to decide whether $SP(n; S) \geq SP(n; R)$ for all $n \geq N(S, R)$, we need to estimate $\prod_{i=1}^T p(n + s_{i,j})$

and $\prod_{i=1}^T p(n + r_{i,j})$ individually for each $1 \leq j \leq M$. In view of Theorem (6.9), estimations of the two factors come into the prominence: computation of the term $\sum_{j=1}^M (\widehat{P}_{n, \vec{s}_j}(w-1) - \widehat{P}_{n, \vec{r}_j}(w-1))$ with $\vec{s}_j := (s_{1,j}, \dots, s_{T,j})$, $\vec{r}_j := (r_{1,j}, \dots, r_{T,j})$, and approximation of the error term.

- (2) Depending on the truncation point w , one can compute the main term $\sum_{j=1}^M (\widehat{P}_{n, \vec{s}_j}(w-1) - \widehat{P}_{n, \vec{r}_j}(w-1))$. But computational complexity will arise in the estimation of the error term because in order to approximate $\widehat{E}(w; \vec{s}_j)$ for each j , one needs to have a good control over $X_{\mathcal{E}}(k)$ for $0 \leq k \leq w-1$. This seems to be difficult as w tends to infinity, growth of $|X_{\mathcal{E}}(k)|$ is exponential.
- (3) For example, in order to prove the higher order Turán inequality for $p(n)$, the minimal choice for w is 10 and consequently, by Theorem 6.9 with appropriate choices for \vec{s} , it follows that

$$4(1 - u_{n-1})(1 - u_n) - (1 - u_n u_{n-1})^2 = \frac{\pi^3}{12\sqrt{6}} \frac{1}{n^{9/2}} + O\left(\frac{1}{n^5}\right).$$

This concludes that $p(n)$ satisfies the higher order Turán inequality for sufficiently large n although due to Chen, Jia, and Wang [6], we know that the inequality holds for all $n \geq 95$. So, from the aspect of error bound computation in order to confirm such inequalities from a certain explicit point onward, our method is inaccessible.

- (4) Last, but not the least, the above discussions naively suggest that for making a decision whether a given inequality for the partition function (of the above type) holds or not, we need to have a full asymptotic expansion for shifted value of the partition function and an explicit computation of the error bound after truncation the expansion at any positive integer w . Recently, Paule, Radu, Schneider, and the author [2] provided an explicit estimation of the error term for the asymptotic expansion of $p(n)$ and following a similar approach, full asymptotic expansion of $p(n-j)$ (with $j \geq 0$) along with an estimation of the error term has been computed in [1].

Acknowledgments

The author sincerely thanks the referee for his/her helpful suggestions and comments. The research was funded by the Austrian Science Fund (FWF): W1214-N15, project DK6.

References

- [1] K. Banerjee, Invariants of the quartic binary form and proofs of Chen's conjectures for partition function inequalities, submitted for publication, 2022. Available at https://www3.risc.jku.at/publications/download/risc_6615/Chen.pdf.
- [2] K. Banerjee, P. Paule, C.S. Radu, C. Schneider, Error bounds for the asymptotic expansion of the partition function, *Rocky Mt. J. Math.* (2023), in press. Available at <https://doi.org/10.48550/arXiv.2209.07887>.
- [3] K. Banerjee, P. Paule, C.S. Radu, W.H. Zeng, New inequalities for $p(n)$ and $\log p(n)$, *Ramanujan J.* 61 (2023) 1295–1338.
- [4] C. Bessenrodt, K. Ono, Maximal multiplicative properties of partitions, *Ann. Comb.* 20 (2016) 59–64.
- [5] W.Y.C. Chen, Recent developments on log-concavity and q -log-concavity of combinatorial polynomials, in: *DMTCS Proceeding of 22nd International Conference on Formal Power Series and Algebraic Combinatorics*, 2010.
- [6] W.Y.C. Chen, D. Jia, L. Wang, Higher order Turán inequalities for the partition function, *Trans. Am. Math. Soc.* 372 (2019) 2143–2165.
- [7] W.Y.C. Chen, L. Wang, G. Xie, Finite differences of the logarithm of the partition function, *Math. Comput.* 85 (2016) 825–847.
- [8] S. DeSalvo, I. Pak, Log-concavity of the partition function, *Ramanujan J.* 38 (2015) 61–73.
- [9] L.E. Dickson, *History of the Theory of Numbers*, vol. 2, Carnegie Institution, Washington, 1920; reprinted: Chelsea, New York, 1952.
- [10] C. Goldbach, *Lettre CXXXI*, <http://eulerarchive.maa.org/correspondence/letters/OO0859.pdf> (18 July, 1750).
- [11] R.L. Graham, D.E. Knuth, O. Patashnik, *Concrete Mathematics*, second edition, Addison Wesley, Reading MA, 1994.
- [12] G.H. Hardy, S. Ramanujan, Asymptotic formulae in combinatory analysis, *Proc. Lond. Math. Soc.* 2 (1918) 75–115.
- [13] D.H. Lehmer, On the remainders and convergence of the series for the partition function, *Trans. Am. Math. Soc.* 46 (1939) 362–373.
- [14] D.H. Lehmer, On the series for the partition function, *Trans. Am. Math. Soc.* 43 (1938) 271–295.
- [15] M. Merca, J. Katriel, A general method for proving the non-trivial linear homogeneous partition inequalities, *Ramanujan J.* 51 (2020) 245–266.
- [16] J.L. Nicolas, Sur les entiers N pour lesquels il y a beaucoup de groupes abéliens d'ordre N , *Ann. Inst. Fourier* 28 (1978) 1–16.
- [17] A.M. Odlyzko, Differences of the partition function, *Acta Arith.* 49 (1988) 237–254.
- [18] H. Rademacher, A convergent series for the partition function $p(n)$, *Proc. Natl. Acad. Sci. USA* 23 (1937) 78.
- [19] H. Rademacher, On the expansion of the partition function in a series, *Ann. Math.* 44 (1943) 416–422.
- [20] H. Rademacher, On the partition function $p(n)$, *Proc. Lond. Math. Soc.* 2 (1938) 241–254.