

## HIGHER ORDER TURÁN INEQUALITIES FOR THE PARTITION FUNCTION

WILLIAM Y. C. CHEN, DENNIS X. Q. JIA, AND LARRY X. W. WANG

ABSTRACT. The Turán inequalities and the higher order Turán inequalities arise in the study of the Maclaurin coefficients of real entire functions in the Laguerre–Pólya class. A sequence  $\{a_n\}_{n \geq 0}$  of real numbers is said to satisfy the Turán inequalities or to be log-concave if for  $n \geq 1$ ,  $a_n^2 - a_{n-1}a_{n+1} \geq 0$ . It is said to satisfy the higher order Turán inequalities if for  $n \geq 1$ ,  $4(a_n^2 - a_{n-1}a_{n+1})(a_{n+1}^2 - a_n a_{n+2}) - (a_n a_{n+1} - a_{n-1} a_{n+2})^2 \geq 0$ . For the partition function  $p(n)$ , DeSalvo and Pak showed that for  $n > 25$ , the sequence  $\{p(n)\}_{n > 25}$  is log-concave, that is,  $p(n)^2 - p(n-1)p(n+1) > 0$  for  $n > 25$ . It was conjectured by the first author that  $p(n)$  satisfies the higher order Turán inequalities for  $n \geq 95$ . In this paper, we prove this conjecture by using the Hardy–Ramanujan–Rademacher formula to derive an upper bound and a lower bound for  $p(n+1)p(n-1)/p(n)^2$ . Consequently, for  $n \geq 95$ , the Jensen polynomials  $p(n-1) + 3p(n)x + 3p(n+1)x^2 + p(n+2)x^3$  have only distinct real zeros. We conjecture that for any positive integer  $m \geq 4$  there exists an integer  $N(m)$  such that for  $n \geq N(m)$ , the Jensen polynomial associated with the sequence  $(p(n), p(n+1), \dots, p(n+m))$  has only real zeros. This conjecture was posed independently by Ono.

### 1. INTRODUCTION

The objective of this paper is to prove the higher order Turán inequalities for the partition function  $p(n)$  when  $n \geq 95$ , as conjectured in [6]. The Turán inequalities and the higher order Turán inequalities are related to the Laguerre–Pólya class of real entire functions [14, 44]. In this paper, a sequence  $\{a_n\}_{n \geq 0}$  always means a sequence of real numbers, and it is said to satisfy the Turán inequalities or to be log-concave if

$$(1.1) \quad a_n^2 - a_{n-1}a_{n+1} \geq 0$$

for  $n \geq 1$ . The inequalities (1.1) are also called the Newton inequalities [8, 11, 35, 46]. We say that a sequence  $\{a_n\}_{n \geq 0}$  satisfies the higher order Turán inequalities or cubic Newton inequalities if for  $n \geq 1$ ,

$$(1.2) \quad 4(a_n^2 - a_{n-1}a_{n+1})(a_{n+1}^2 - a_n a_{n+2}) - (a_n a_{n+1} - a_{n-1} a_{n+2})^2 \geq 0;$$

see [14, 35, 41]. It is worth mentioning the double Turán inequalities and the higher order iterated Turán inequalities introduced by Csordas [10]; see also [24]. Given a sequence  $\{a_n\}_{n \geq 0}$ , for  $n \geq 1$ , let

$$T_1(n) = a_n^2 - a_{n-1}a_{n+1},$$

---

Received by the editors August 5, 2017, and, in revised form, September 19, 2018.

2010 *Mathematics Subject Classification*. Primary 05A20, 11P82.

*Key words and phrases*. Partition function, log-concavity, higher order Turán inequalities, Hardy–Ramanujan–Rademacher formula, Jensen polynomials.

This work was supported by the National Science Foundation of China.

and for  $k \geq 2$  and  $n \geq k$ , let

$$T_k(n) = T_{k-1}(n)^2 - T_{k-1}(n-1)T_{k-1}(n+1).$$

A sequence  $\{a_n\}_{n \geq 0}$  is said to satisfy the double Turán inequalities if  $T_1(n) \geq 0$  for  $n \geq 1$  and  $T_2(n) \geq 0$  for  $n \geq 2$ . In general,  $\{a_n\}_{n \geq 0}$  is said to satisfy the  $l$ th order iterated Turán inequalities if, for  $1 \leq k \leq l$  and  $n \geq k$ , we have  $T_k(n) \geq 0$ . It should be noted that the above notion of higher order iterated Turán inequalities coincides with the notion of the higher order log-concavity introduced by Moll [34]; see also [3, 4]. In the terminology of Moll, a sequence  $\{a_n\}_{n \geq 0}$  satisfying the  $l$ th order iterated Turán inequalities is called  $l$ -log-concave. A sequence  $\{a_n\}_{n \geq 0}$  is said to be infinitely log-concave if it is  $l$ -log-concave for any  $l \geq 1$ .

It was conjectured by Boros and Moll [3] that for each  $n \geq 0$ , the sequence of binomial coefficients  $\{\binom{n}{k}\}_{k=0}^n$  is infinitely log-concave. A stronger conjecture was proposed independently by Stanley [4], Fisk [17], and McNamara and Sagan [32], which states that if  $a_0 + a_1x + \dots + a_nx^n$  is a real-rooted polynomial with nonnegative coefficients, then so is the polynomial  $b_0 + b_1x + \dots + b_nx^n$ , where  $b_0 = a_0$ ,  $b_k = a_k^2 - a_{k+1}a_{k-1}$  for  $1 \leq k \leq n-1$ , and  $b_n = a_n$ . This conjecture has been proved by Brändén [4].

A real entire function

$$(1.3) \quad \psi(x) = \sum_{k=0}^{\infty} \gamma_k \frac{x^k}{k!}$$

is said to be in the Laguerre–Pólya class, denoted  $\psi(x) \in \mathcal{LP}$ , if it can be represented in the form

$$\psi(x) = cx^m e^{-\alpha x^2 + \beta x} \prod_{k=1}^{\infty} (1 + x/x_k) e^{-x/x_k},$$

where  $c, \beta, x_k$  are real numbers,  $\alpha \geq 0$ ,  $m$  is a nonnegative integer, and  $\sum x_k^{-2} < \infty$ . These functions are the only ones which are uniform limits of polynomials whose zeros are real. We refer to [29] and [42] for the background on the theory of the  $\mathcal{LP}$  class.

Jensen [22] proved that a real entire function  $\psi(x)$  belongs to the  $\mathcal{LP}$  class if and only if for any positive integer  $m$ , the  $m$ th associated Jensen polynomial

$$(1.4) \quad g_m(x) = \sum_{k=0}^m \binom{m}{k} \gamma_k x^k$$

has only real zeros. More properties of the Jensen polynomials can be found in [8, 11, 12].

Pólya and Schur [39] also obtained the above result based on multiplier sequences of the second kind. A real sequence  $\{\gamma_k\}_{k \geq 0}$  is called a multiplier sequence of the second kind if for any nonnegative integer  $m$  and every real polynomial  $\sum_{k=0}^m a_k x^k$  with only real zeros of the same sign, the polynomial  $\sum_{k=0}^m a_k \gamma_k x^k$  has only real zeros. Pólya and Schur [39] proved that each multiplier sequence of the second kind satisfies the Turán inequalities. Moreover, they showed that a real entire function  $\psi(x)$  belongs to the  $\mathcal{LP}$  class if and only if the sequence of its Maclaurin coefficients is a multiplier sequence of the second kind. This implies that the Maclaurin coefficients of  $\psi(x)$  in the  $\mathcal{LP}$  class satisfy the Turán inequalities

$$(1.5) \quad \gamma_k^2 - \gamma_{k-1}\gamma_{k+1} \geq 0$$

for  $k \geq 1$ . In fact, (1.5) is a consequence of another property of the  $\mathcal{LP}$  class due to Pólya and Schur [39]: Let  $\psi(x)$  be a real entire function in the  $\mathcal{LP}$  class. For any nonnegative integer  $n$ , the  $n$ th derivative  $\psi^{(n)}(x)$  of  $\psi(x)$  also belongs to the  $\mathcal{LP}$  class. It is readily seen that the  $m$ th Jensen polynomial associated with  $\psi^{(n)}(x)$  is

$$(1.6) \quad g_{m,n}(x) = \sum_{k=0}^m \binom{m}{k} \gamma_{k+n} x^k,$$

and hence it has only real zeros for any nonnegative integers  $m$  and  $n$ . In particular, taking  $m = 2$ , for any nonnegative integer  $n$ , the real-rootedness of  $g_{2,n}(x)$  implies the inequality (1.5), that is, the discriminant  $4(\gamma_{n+1}^2 - \gamma_n \gamma_{n+2})$  is nonnegative.

Dimitrov [14] observed that for a real entire function  $\psi(x)$  in the  $\mathcal{LP}$  class, the Maclaurin coefficients satisfy the higher order Turán inequalities

$$(1.7) \quad 4(\gamma_k^2 - \gamma_{k-1} \gamma_{k+1})(\gamma_{k+1}^2 - \gamma_k \gamma_{k+2}) - (\gamma_k \gamma_{k+1} - \gamma_{k-1} \gamma_{k+2})^2 \geq 0$$

for  $k \geq 1$ . This fact follows from a theorem of Mařík [31] stating that if the real polynomial

$$(1.8) \quad \sum_{k=0}^m \binom{m}{k} a_k x^k$$

of degree  $m \geq 3$  has only real zeros, then  $a_0, a_1, \dots, a_m$  satisfy the higher order Turán inequalities.

As noted in [6], for  $k = 1$ , the polynomial in (1.7) coincides with an invariant

$$I(a_0, a_1, a_2, a_3) = 3a_1^2 a_2^2 - 4a_1^3 a_3 - 4a_0 a_2^3 - a_0^2 a_3^2 + 6a_0 a_1 a_2 a_3$$

of the cubic binary form

$$(1.9) \quad a_3 x^3 + 3a_2 x^2 y + 3a_1 x y^2 + a_0 y^3.$$

In other words, the above invariant  $I(a_0, a_1, a_2, a_3)$  can be rewritten as

$$(1.10) \quad I(a_0, a_1, a_2, a_3) = 4(a_1^2 - a_0 a_2)(a_2^2 - a_1 a_3) - (a_1 a_2 - a_0 a_3)^2.$$

We refer to Hilbert [20], Kung and Rota [25], and Sturmfels [43] for the background on the invariant theory of binary forms. Notice that  $27I(a_0, a_1, a_2, a_3)$  is the discriminant of the cubic polynomial  $a_3 x^3 + 3a_2 x^2 + 3a_1 x + a_0$  [33]. For a real cubic polynomial, the discriminant is positive if and only if the three zeros are real and distinct. In general, for a real polynomial of degree greater than or equal to four, the discriminant is positive if and only if the number of nonreal roots is a multiple of four. More properties about discriminants can be found in [21] and [33].

Recall that for a real entire function  $\psi(x)$  in the  $\mathcal{LP}$  class, its  $n$ th derivative  $\psi^{(n)}(x)$  is also a real entire function in the  $\mathcal{LP}$  class. Thus the real-rootedness of the cubic Jensen polynomial  $g_{3,n}(x)$  associated with  $\psi^{(n)}(x)$  implies the higher order Turán inequalities (1.7) of Dimitrov, that is, the discriminant  $27I(\gamma_n, \gamma_{n+1}, \gamma_{n+2}, \gamma_{n+3})$  is nonnegative.

Let  $f(x)$  be a real polynomial with degree  $m$ . If  $x_1, x_2, \dots, x_m$  are the roots of  $f(x)$ , let  $S_k = \sum_{i=1}^m x_i^k$  denote the  $k$ th Newton sums of  $f(x)$  for  $k \geq 1$  and  $S_0 = m$ . Explicit expressions for these power sums via the coefficients of  $f(x)$  are given by the Waring formula [30]. The Hermite matrix of  $f(x)$  is a symmetric  $m \times m$  matrix, defined as

$$\mathcal{H}_m(f(x)) = (S_{i+j-2})_{i,j=1,2,\dots,m}.$$

It is a Hankel matrix whose entries are polynomials in the coefficients of  $f(x)$ . We write  $D_k(f(x))$  for the leading principal minor of  $\mathcal{H}_m(f(x))$  of order  $k$ . A theorem of Hermite [37] says that all the zeros of  $f(x)$  are real if and only if all the leading principal minors of  $\mathcal{H}_m(f(x))$  are nonnegative. By this theorem, one can state that many more (higher Turán) inequalities would need to be true in order for the higher degree Jensen polynomials to have only real zeros. For example, consider the Jensen polynomials  $g_{4,k}(x)$  for  $k \geq 0$ , that is,

$$g_{4,k}(x) = \gamma_k + 4\gamma_{k+1}x + 6\gamma_{k+2}x^2 + 4\gamma_{k+3}x^3 + \gamma_{k+4}x^4.$$

It can be calculated that for  $k \geq 0$ ,

$$\begin{aligned} D_1(g_{4,k}(x)) &= 4, & D_2(g_{4,k}(x)) &= \frac{48(\gamma_{k+3}^2 - \gamma_{k+2}\gamma_{k+4})}{\gamma_{k+4}^2}, \\ D_3(g_{4,k}(x)) &= \frac{192A}{\gamma_{k+4}^4}, & \det(\mathcal{H}_4(g_{4,k}(x))) &= \frac{256(B^3 - 27C^2)}{\gamma_{k+4}^6}, \end{aligned}$$

where

$$\begin{aligned} A &= 6\gamma_{k+2}^2\gamma_{k+3}^2 + \gamma_k\gamma_{k+2}\gamma_{k+4}^2 + 14\gamma_{k+1}\gamma_{k+2}\gamma_{k+3}\gamma_{k+4} - 9\gamma_{k+4}\gamma_{k+2}^3 \\ &\quad - 8\gamma_{k+1}\gamma_{k+3}^3 - 3\gamma_{k+1}^2\gamma_{k+4}^2 - \gamma_k\gamma_{k+4}\gamma_{k+3}^2, \\ B &= 3\gamma_{k+2}^2 - 4\gamma_{k+1}\gamma_{k+3} + \gamma_k\gamma_{k+4}, \\ C &= \gamma_{k+2}^3 - 2\gamma_{k+1}\gamma_{k+2}\gamma_{k+3} - \gamma_k\gamma_{k+2}\gamma_{k+4} + \gamma_k\gamma_{k+3}^2 + \gamma_{k+1}^2\gamma_{k+4}. \end{aligned}$$

Notice that  $B, C$  and  $B^3 - 27C^2$  are actually the invariants of the quartic binary form with the same coefficients as  $g_{4,k}(x)$  [16, 20] and  $256(B^3 - 27C^2)$  is the discriminant of  $g_{4,k}(x)$ . By Hermite's theorem, one of the necessary and sufficient conditions for  $g_{4,k}(x)$  to have only real zeros is that all the leading principal minors of  $\mathcal{H}_4(g_{4,k}(x))$  are nonnegative, that is, for  $k \geq 0$ ,

$$(1.11) \quad \gamma_{k+3}^2 - \gamma_{k+2}\gamma_{k+4} \geq 0, \quad A \geq 0, \quad B^3 - 27C^2 \geq 0.$$

Real entire functions in the  $\mathcal{LP}$  class with nonnegative Maclaurin coefficients have also received much attention. Aissen, Schoenberg and Whitney [1] proved that if  $\psi(x)$  is a real entire function in the  $\mathcal{LP}$  class with nonnegative Maclaurin coefficients, then the sequence  $\{\gamma_k/k!\}$  associated with  $\psi(x)$  forms a Pólya frequency sequence. An infinite sequence  $\{a_n\}_{n \geq 0}$  of nonnegative numbers is called a Pólya frequency sequence (or a  $PF$ -sequence) if the matrix  $(a_{i-j})_{i,j \geq 0}$  is a totally positive matrix, where  $a_n = 0$  if  $n < 0$ ; that is, all minors of  $(a_{i-j})_{i,j \geq 0}$  have nonnegative determinants. More properties of totally positive matrices and  $PF$ -sequences can be found in [9, 23].

The  $\mathcal{LP}$  class is closely related to the Riemann hypothesis. Let  $\zeta$  denote the Riemann zeta-function, and let  $\Gamma$  be the gamma-function. The Riemann  $\xi$ -function is defined by

$$(1.12) \quad \xi(iz) = \frac{1}{2} \left( z^2 - \frac{1}{4} \right) \pi^{-z/2-1/4} \Gamma \left( \frac{z}{2} + \frac{1}{4} \right) \zeta \left( z + \frac{1}{2} \right);$$

see, for example, Boas [2]. It is well known that the Riemann  $\xi$ -function is an entire function of order 1 and can be represented in the following form [38]:

$$(1.13) \quad \frac{1}{8}\xi\left(\frac{x}{2}\right) = \sum_{k=0}^{\infty} (-1)^k \hat{b}_k \frac{x^{2k}}{(2k)!}, \quad k = 0, 1, 2, \dots,$$

where

$$\hat{b}_k = \int_0^{\infty} t^{2k} \Phi(t) dt \quad \text{and} \quad \Phi(t) = \sum_{n=0}^{\infty} (2n^4 \pi^2 e^{9t} - 3n^2 \pi e^{5t}) \exp(-n^2 \pi e^{4t}).$$

Setting  $z = -x^2$  in (1.13), we are led to an entire function of order 1/2, denoted  $\xi_1(z)$ , that is,

$$(1.14) \quad \xi_1(z) = \sum_{k=0}^{\infty} \hat{\gamma}_k \frac{z^k}{k!},$$

where

$$\hat{\gamma}_k = \frac{k!}{(2k)!} \hat{b}_k, \quad k = 0, 1, 2, \dots$$

Thus, the Riemann hypothesis holds if and only if  $\xi_1(z)$  belongs to the  $\mathcal{LP}$  class. We note that  $\xi_1(z)$  has no positive zeros since  $\hat{\gamma}_k$  is positive for  $k \geq 0$ . For a real entire function  $\psi(x)$  as defined in (1.3), it is well known that  $\psi(x)$  belongs to  $\mathcal{LP}$  if and only if the Jensen polynomials  $g_{m,n}(x)$  in (1.6) have only real zeros [12, 37]. Let  $\hat{g}_{m,n}(x)$  denote the Jensen polynomials associated with  $\xi_1(z)$ , that is,

$$\hat{g}_{m,n}(x) = \sum_{k=0}^m \binom{m}{k} \hat{\gamma}_{k+n} x^k, \quad m, n = 0, 1, 2, \dots,$$

then the Riemann hypothesis is equivalent to the statement that  $\hat{g}_{m,n}(x)$  has only real zeros for  $m \geq 1$  and  $n \geq 0$ . For  $m = 2$  and  $n \geq 0$ ,  $\hat{g}_{2,n}(x)$  has only real zeros if and only if the discriminant of  $\hat{g}_{2,n}(x)$  is nonnegative, that is, for  $k \geq 1$ ,

$$(1.15) \quad \hat{\gamma}_k^2 - \hat{\gamma}_{k-1} \hat{\gamma}_{k+1} \geq 0.$$

The above inequalities (1.15) were proved by Csordas, Norfolk, and Varga [11]. This shows that for each  $n \geq 0$ ,  $\hat{g}_{2,n}(x)$  has only real zeros. For  $m = 3$  and  $n \geq 0$ , the real-rootness of  $\hat{g}_{3,n}(x)$  can be deduced from the higher order Turán inequalities (1.7), that is, for  $k \geq 1$ ,

$$4(\hat{\gamma}_k^2 - \hat{\gamma}_{k-1} \hat{\gamma}_{k+1})(\hat{\gamma}_{k+1}^2 - \hat{\gamma}_k \hat{\gamma}_{k+2}) - (\hat{\gamma}_k \hat{\gamma}_{k+1} - \hat{\gamma}_{k-1} \hat{\gamma}_{k+2})^2 \geq 0,$$

as proved by Dimitrov and Lucas [15]. Recently, Griffin, Ono, Rolen, and Zagier [18] proved that for each  $m \geq 1$ ,  $\hat{g}_{m,n}(x)$  has only real zeros with at most finitely many exceptions  $n$ .

Let us now turn to the partition function. A partition of a positive integer  $n$  is a nonincreasing sequence  $(\lambda_1, \lambda_2, \dots, \lambda_r)$  of positive integers such that  $\lambda_1 + \lambda_2 + \dots + \lambda_r = n$ . Let  $p(n)$  denote the number of partitions of  $n$ . DeSalvo and Pak [13] proved the log-concavity of the partition function for  $n > 25$  as well as the following inequality as conjectured in [5]: for  $n \geq 2$ ,

$$(1.16) \quad \frac{p(n-1)}{p(n)} \left(1 + \frac{1}{n}\right) > \frac{p(n)}{p(n+1)}.$$

DaSalvo and Pak also conjectured that for  $n \geq 45$ ,

$$(1.17) \quad \frac{p(n-1)}{p(n)} \left( 1 + \frac{\pi}{\sqrt{24}n^{3/2}} \right) > \frac{p(n)}{p(n+1)}.$$

Chen, Wang, and Xie [7] gave an affirmative answer to this conjecture.

It was conjectured in [6] that for large  $n$ , the partition function  $p(n)$  satisfies many inequalities pertaining to invariants of a binary form. Here we mention two of them.

**Conjecture 1.1.** *For  $n \geq 95$ , the higher order Turán inequalities*

$$(1.18) \quad 4(a_n^2 - a_{n-1}a_{n+1})(a_{n+1}^2 - a_n a_{n+2}) - (a_n a_{n+1} - a_{n-1} a_{n+2})^2 \geq 0$$

hold for  $a_n = p(n)$ .

The following conjecture is a higher order analogue of (1.17).

**Conjecture 1.2.** *Let*

$$(1.19) \quad u_n = \frac{p(n+1)p(n-1)}{p(n)^2}.$$

For  $n \geq 2$ ,

$$4(1 - u_n)(1 - u_{n+1}) < \left( 1 + \frac{\pi}{\sqrt{24}n^{3/2}} \right) (1 - u_n u_{n+1})^2.$$

The objective of this paper is to prove Conjecture 1.1. In fact, we shall prove the following equivalent form.

**Theorem 1.3.** *Let  $u_n$  be defined as in (1.19). For  $n \geq 95$ ,*

$$(1.20) \quad 4(1 - u_n)(1 - u_{n+1}) - (1 - u_n u_{n+1})^2 > 0.$$

The above theorem can be restated as follows.

**Theorem 1.4.** *For any  $n \geq 95$ , the cubic polynomial*

$$p(n-1) + 3p(n)x + 3p(n+1)x^2 + p(n+2)x^3$$

has three distinct real zeros.

In general, we propose the following conjecture.

**Conjecture 1.5.** *For any positive integer  $m \geq 4$ , there exists a positive integer  $N(m)$  such that for any  $n \geq N(m)$ , the Jensen polynomial*

$$\sum_{k=0}^m \binom{m}{k} p(k+n)x^k$$

has only real zeros.

The above conjecture was proposed independently by Ono [36]. For fixed  $m$  and large  $n$ , Griffin, Ono, Rolin, and Zagier [18] proved that this conjecture is true. In fact, they showed that for suitable entire functions and certain sequences, the associated Jensen polynomials have only real and distinct zeros with at most finitely many exceptions. To be more precise, they defined the normalized Jensen polynomials by changing the variable of the Jensen polynomials and proved that for large  $n$ , the normalized Jensen polynomials are small perturbations of the Hermite polynomials  $H_m(x)$ . Since all the roots of  $H_m(x)$  are real and distinct [45], the real

parts of the roots of the normalized Jensen polynomials are distinct for large  $n$ , which implies that all the roots are real and distinct. According to the definition of the normalized Jensen polynomials, it is easy to see that for large  $n$ , all the roots of such Jensen polynomials are also real and distinct.

Assume that  $N(m)$  is the minimum value in Conjecture 1.5. Larson and Wagner [26] showed that  $N(3) = 94$ ,  $N(4) = 206$ , and  $N(5) = 381$ , and that  $N(m) \leq (3m)^{24m}(50m)^{3m^2}$  for  $m \geq 1$ . They also gave a proof of Conjecture 1.2.

## 2. BOUNDING $u_n$

In this section, we give an upper bound and a lower bound for

$$u_n = \frac{p(n+1)p(n-1)}{p(n)^2},$$

as defined in (1.19). DeSalvo and Pak [13] proved that for  $n > 25$ ,

$$1 - \frac{1}{n+1} < u_n < 1.$$

On the other hand, Chen, Wang and Xie [7] showed that for  $n \geq 45$ ,

$$1 - \frac{\pi}{\sqrt{24n^{3/2} + \pi}} < u_n.$$

Nevertheless, the above bounds for  $u_n$  are not sharp enough for the purpose of proving Theorem 1.3. To state our bounds for  $u_n$ , we adopt the following notation as used in [28]:

$$(2.1) \quad \mu(n) = \frac{\pi}{6} \sqrt{24n - 1}.$$

For convenience, let

$$(2.2) \quad x = \mu(n-1), \quad y = \mu(n), \quad z = \mu(n+1), \quad w = \mu(n+2).$$

Define

$$(2.3) \quad f(n) = e^{x-2y+z} \frac{(x^{10} - x^9 - 1) y^{24} (z^{10} - z^9 - 1)}{x^{12} (y^{10} - y^9 + 1)^2 z^{12}},$$

$$(2.4) \quad g(n) = e^{x-2y+z} \frac{(x^{10} - x^9 + 1) y^{24} (z^{10} - z^9 + 1)}{x^{12} (y^{10} - y^9 - 1)^2 z^{12}}.$$

Then we have the following bounds for  $u_n$ .

**Theorem 2.1.** *For  $n \geq 1207$ ,*

$$(2.5) \quad f(n) < u_n < g(n).$$

To prove Theorem 2.1, we need the following upper bound and lower bound for  $p(n)$ .

**Lemma 2.2.** *Let*

$$B_1(n) = \frac{\sqrt{12}e^{\mu(n)}}{24n - 1} \left( 1 - \frac{1}{\mu(n)} - \frac{1}{\mu(n)^{10}} \right),$$

$$B_2(n) = \frac{\sqrt{12}e^{\mu(n)}}{24n - 1} \left( 1 - \frac{1}{\mu(n)} + \frac{1}{\mu(n)^{10}} \right),$$

then for  $n \geq 1206$ ,

$$(2.6) \quad B_1(n) < p(n) < B_2(n).$$

The proof of Lemma 2.2 relies on the Hardy–Ramanujan–Rademacher formula [19, 40] for  $p(n)$  as well as Lehmer’s error bound for the remainder of this formula. The Hardy–Ramanujan–Rademacher formula reads

$$(2.7) \quad p(n) = \frac{\sqrt{12}}{24n - 1} \sum_{k=1}^N \frac{A_k(n)}{\sqrt{k}} \left[ \left(1 - \frac{k}{\mu(n)}\right) e^{\mu(n)/k} + \left(1 + \frac{k}{\mu(n)}\right) e^{-\mu(n)/k} \right] + R_2(n, N),$$

where  $A_k(n)$  is an arithmetic function and  $R_2(n, N)$  is the remainder term; see, for example, Rademacher [40]. Lehmer [27, 28] gave the error bound

$$(2.8) \quad |R_2(n, N)| < \frac{\pi^2 N^{-2/3}}{\sqrt{3}} \left[ \left(\frac{N}{\mu(n)}\right)^3 \sinh \frac{\mu(n)}{N} + \frac{1}{6} - \left(\frac{N}{\mu(n)}\right)^2 \right],$$

which is valid for all positive integers  $n$  and  $N$ .

*Proof of Lemma 2.2.* Consider the Hardy–Ramanujan–Rademacher formula (2.7) for  $N = 2$ , and note that  $A_1(n) = 1$  and  $A_2(n) = (-1)^n$  for any positive integer  $n$ . Hence (2.7) can be rewritten as

$$(2.9) \quad p(n) = \frac{\sqrt{12}e^{\mu(n)}}{24n - 1} \left(1 - \frac{1}{\mu(n)} + T(n)\right),$$

where

$$(2.10) \quad T(n) = \frac{(-1)^n}{\sqrt{2}} \left( \left(1 - \frac{2}{\mu(n)}\right) e^{-\mu(n)/2} + \left(1 + \frac{2}{\mu(n)}\right) e^{-3\mu(n)/2} \right) + \left(1 + \frac{1}{\mu(n)}\right) e^{-2\mu(n)} + \frac{(24n - 1)R_2(n, 2)}{\sqrt{12}e^{\mu(n)}}.$$

In order to prove (2.6), we proceed to use Lehmer’s error bound in (2.8) to show that for  $n > 1520$ ,

$$(2.11) \quad |T(n)| < \frac{1}{\mu(n)^{10}}.$$

Let us begin with the first three terms in (2.10). Evidently, for  $n \geq 1$ ,

$$0 < \frac{1}{\mu(n)} < \frac{1}{2},$$

so that

$$(2.12) \quad \left(1 - \frac{2}{\mu(n)}\right) e^{-\mu(n)/2} < e^{-\mu(n)/2},$$

$$(2.13) \quad \left(1 + \frac{2}{\mu(n)}\right) e^{-3\mu(n)/2} < 2e^{-3\mu(n)/2},$$

$$(2.14) \quad \left(1 + \frac{1}{\mu(n)}\right) e^{-2\mu(n)} < 2e^{-2\mu(n)}.$$

As for the last term in (2.10), we claim that for  $n > 350$ ,

$$(2.15) \quad \left| \frac{(24n - 1)R_2(n, 2)}{\sqrt{12}e^{\mu(n)}} \right| < e^{-\mu(n)/2}.$$



Applying (2.8) with  $N = 2$ , we obtain that

$$\begin{aligned}
 (2.16) \quad & \left| \frac{(24n - 1)R_2(n, 2)}{\sqrt{12}e^{\mu(n)}} \right| = \left| \frac{36\mu(n)^2 R_2(n, 2)}{\sqrt{12}\pi^2 e^{\mu(n)}} \right| \\
 & < \frac{\mu(n)^2 e^{-\mu(n)}}{2^{2/3}} + \frac{12\sqrt[3]{2}e^{-\mu(n)/2}}{\mu(n)} - \frac{12\sqrt[3]{2}e^{-3\mu(n)/2}}{\mu(n)} - 12\sqrt[3]{2}e^{-\mu(n)} \\
 & < \frac{\mu(n)^2 e^{-\mu(n)}}{2^{2/3}} + \frac{12\sqrt[3]{2}e^{-\mu(n)/2}}{\mu(n)} \\
 & < \frac{24e^{-\mu(n)/2}}{\mu(n)} + \mu(n)^2 e^{-\mu(n)}.
 \end{aligned}$$

To bound the first term in (2.16), we see that for  $n > 350$ ,

$$(2.17) \quad \frac{24e^{-\mu(n)/2}}{\mu(n)} < \frac{e^{-\mu(n)/2}}{2},$$

since

$$(2.18) \quad \mu(n) = \frac{\pi}{6}\sqrt{24n - 1} > 48$$

holds for  $n > 350$ . Concerning the second term in (2.16), it will be shown that for  $n > 22$ ,

$$(2.19) \quad \mu(n)^2 e^{-\mu(n)} < \frac{e^{-\mu(n)/2}}{2},$$

which can be rewritten as

$$(2.20) \quad \frac{e^{\mu(n)/4}}{\mu(n)/4} > 4\sqrt{2}.$$

Let

$$(2.21) \quad F(t) = \frac{e^t}{t}.$$

Since  $F'(t) = e^t(t - 1)/t^2 > 0$  for  $t > 1$ ,  $F(t)$  is increasing for  $t > 1$ . Thus,

$$F\left(\frac{\mu(n)}{4}\right) = \frac{e^{\mu(n)/4}}{\mu(n)/4} > F(3) = \frac{e^3}{3} > 4\sqrt{2}.$$

Here we have used the fact that for  $n > 22$ ,  $\mu(n)/4 > 3$ . This proves (2.20). Applying the estimates (2.17) and (2.19) to (2.16), we reach (2.15).

Taking all the above estimates into account, we deduce that for  $n > 350$ ,

$$(2.22) \quad |T(n)| < 6e^{-\mu(n)/2}.$$

To obtain (2.11), we only have to show that for  $n > 1520$ ,

$$(2.23) \quad 6e^{-\mu(n)/2} < \frac{1}{\mu(n)^{10}},$$

which can be recast as

$$(2.24) \quad \frac{e^{\mu(n)/20}}{\mu(n)/20} > 20^{10}\sqrt[10]{6}.$$

Since  $\mu(n)/20 > 5$  for  $n > 1520$ , by the monotone property of  $F(t)$ , we deduce that for  $n > 1520$ ,

$$F\left(\frac{\mu(n)}{20}\right) = \frac{e^{\mu(n)/20}}{\mu(n)/20} > F(5) = \frac{e^5}{5} > 20^{10}\sqrt[10]{6},$$

as asserted by (2.24). Thus (2.11) follows from (2.22) and (2.23). In other words, for  $n > 1520$ ,

$$(2.25) \quad -\frac{1}{\mu(n)^{10}} < T(n) < \frac{1}{\mu(n)^{10}}.$$

Substituting (2.9) into (2.25), we see that (2.6) holds for  $n > 1520$ . It is routine to check that (2.6) is true for  $1206 \leq n \leq 1520$ , and hence the proof is complete.  $\square$

We are now ready to prove Theorem 2.1.

*Proof of Theorem 2.1.* Since  $B_1(n)$  and  $B_2(n)$  are all positive for  $n \geq 1$ , using the bounds for  $p(n)$  in (2.6), we find that for  $n \geq 1207$ ,

$$\frac{B_1(n-1)B_1(n+1)}{B_2(n)^2} < \frac{p(n-1)p(n+1)}{p(n)^2} < \frac{B_2(n-1)B_2(n+1)}{B_1(n)^2}.$$

This proves (2.5).  $\square$

### 3. AN INEQUALITY ON $f(n)$ AND $g(n)$

In this section, we establish an inequality between  $f(n)$  and  $g(n+1)$  which will be used in the proof of Theorem 1.3.

**Theorem 3.1.** For  $n \geq 2$ ,

$$(3.1) \quad g(n+1) < f(n) + \frac{110}{\mu(n-1)^5}.$$

*Proof.* Let  $\mu(n)$  be defined as in (2.1), that is,

$$\mu(n) = \frac{\pi\sqrt{24n-1}}{6},$$

and let

$$x = \mu(n-1), \quad y = \mu(n), \quad z = \mu(n+1), \quad w = \mu(n+2),$$

as defined in (2.2).

Since  $x \geq 0$  for  $n \geq 2$ , we proceed to show that for  $n \geq 2$ ,

$$f(n)x^5 - g(n+1)x^5 + 110 > 0.$$

Let

$$(3.2) \quad \alpha(t) = t^{10} - t^9 + 1, \quad \beta(t) = t^{10} - t^9 - 1.$$

By the definitions of  $f(n)$  and  $g(n)$  as given in (2.3) and (2.4), we obtain that

$$(3.3) \quad f(n)x^5 - g(n+1)x^5 + 110 = \frac{-e^{w+y-2z}t_1 + e^{z+x-2y}t_2 + 110t_3}{t_3},$$

where

$$(3.4) \quad t_1 = x^{12}z^{36}\alpha(y)^3\alpha(w),$$

$$(3.5) \quad t_2 = y^{36}w^{12}\beta(x)\beta(z)^3,$$

$$(3.6) \quad t_3 = x^7y^{12}z^{12}w^{12}\alpha(y)^2\beta(z)^2.$$

Since  $t_3 > 0$  for  $n \geq 2$ , (3.1) is equivalent to

$$(3.7) \quad -e^{w+y-2z}t_1 + e^{z+x-2y}t_2 + 110t_3 > 0,$$

for  $n \geq 2$ . To verify (3.7), we shall estimate  $t_1, t_2, t_3, e^{w+y-2z}$ , and  $e^{x-2y+z}$  in terms of  $x$ . Noting that for  $n \geq 2$ ,

$$(3.8) \quad y = \sqrt{x^2 + \frac{2\pi^2}{3}}, \quad z = \sqrt{x^2 + \frac{4\pi^2}{3}}, \quad w = \sqrt{x^2 + 2\pi^2},$$

we have

$$y = x + \frac{\pi^2}{3x} - \frac{\pi^4}{18x^3} + \frac{\pi^6}{54x^5} - \frac{5\pi^8}{648x^7} + \frac{7\pi^{10}}{1944x^9} - \frac{7\pi^{12}}{3888x^{11}} + o\left(\frac{1}{x^{12}}\right),$$

$$z = x + \frac{2\pi^2}{3x} - \frac{2\pi^4}{9x^3} + \frac{4\pi^6}{27x^5} - \frac{10\pi^8}{81x^7} + \frac{28\pi^{10}}{243x^9} - \frac{28\pi^{12}}{243x^{11}} + o\left(\frac{1}{x^{12}}\right),$$

$$w = x + \frac{\pi^2}{x} - \frac{\pi^4}{2x^3} + \frac{\pi^6}{2x^5} - \frac{5\pi^8}{8x^7} + \frac{7\pi^{10}}{8x^9} - \frac{21\pi^{12}}{16x^{11}} + o\left(\frac{1}{x^{12}}\right).$$

It is readily checked that for  $x \geq 4$ ,

$$(3.9) \quad y_1 < y < y_2,$$

$$(3.10) \quad z_1 < z < z_2,$$

$$(3.11) \quad w_1 < w < w_2,$$

where

$$(3.12) \quad y_1 = x + \frac{\pi^2}{3x} - \frac{\pi^4}{18x^3} + \frac{\pi^6}{54x^5} - \frac{5\pi^8}{648x^7} + \frac{7\pi^{10}}{1944x^9} - \frac{7\pi^{12}}{3888x^{11}},$$

$$(3.13) \quad y_2 = x + \frac{\pi^2}{3x} - \frac{\pi^4}{18x^3} + \frac{\pi^6}{54x^5} - \frac{5\pi^8}{648x^7} + \frac{7\pi^{10}}{1944x^9},$$

$$(3.14) \quad z_1 = x + \frac{2\pi^2}{3x} - \frac{2\pi^4}{9x^3} + \frac{4\pi^6}{27x^5} - \frac{10\pi^8}{81x^7} + \frac{28\pi^{10}}{243x^9} - \frac{28\pi^{12}}{243x^{11}},$$

$$(3.15) \quad z_2 = x + \frac{2\pi^2}{3x} - \frac{2\pi^4}{9x^3} + \frac{4\pi^6}{27x^5} - \frac{10\pi^8}{81x^7} + \frac{28\pi^{10}}{243x^9},$$

$$(3.16) \quad w_1 = x + \frac{\pi^2}{x} - \frac{\pi^4}{2x^3} + \frac{\pi^6}{2x^5} - \frac{5\pi^8}{8x^7} + \frac{7\pi^{10}}{8x^9} - \frac{21\pi^{12}}{16x^{11}},$$

$$(3.17) \quad w_2 = x + \frac{\pi^2}{x} - \frac{\pi^4}{2x^3} + \frac{\pi^6}{2x^5} - \frac{5\pi^8}{8x^7} + \frac{7\pi^{10}}{8x^9}.$$

With these bounds of  $y, z$ , and  $w$  in (3.9), (3.10), and (3.11), we are now in a position to estimate  $t_1, t_2, t_3, e^{w+y-2z}$ , and  $e^{x-2y+z}$  in terms of  $x$ .

First, we consider  $t_1, t_2$ , and  $t_3$ . By the definition of  $\alpha(t)$ , we have

$$\alpha(w) = w^{10} - w^9 + 1.$$

Noting that  $w^9 = (x^2 + 2\pi^2)^4 \sqrt{x^2 + 2\pi^2}$ , which involves a radical, to give a feasible estimate for  $w^9$  without a radical, we may make use of (3.11) to deduce that for  $x \geq 4$ ,

$$w_1 w^8 < w^9 < w_2 w^8.$$

Let

$$\eta_1 = w^{10} - w_1w^8 + 1$$

so that for  $x \geq 4$ ,

$$(3.18) \quad \alpha(w) < \eta_1.$$

Similarly, set

$$\eta_2 = y^{30} - 3y_1y^{28} + 3y^{28} - y_1y^{26} + 3y^{20} - 6y_1y^{18} + 3y^{18} + 3y^{10} - 3y_1y^8 + 1,$$

$$\eta_3 = z^{30} - 3z_2z^{28} + 3z^{28} - z_2z^{26} - 3z^{20} + 6z_1z^{18} - 3z^{18} + 3z^{10} - 3z_2z^8 - 1,$$

$$\eta_4 = y^{20} - 2y_2y^{18} + y^{18} + 2y^{10} - 2y_2y^8 + 1,$$

$$\eta_5 = z^{20} - 2z_2z^{18} + z^{18} - 2z^{10} + 2z_1z^8 + 1.$$

Then we have for  $x \geq 4$ ,

$$(3.19) \quad \alpha(y)^3 < \eta_2, \quad \beta(z)^3 > \eta_3, \quad \alpha(y)^2 > \eta_4, \quad \beta(z)^2 > \eta_5.$$

Employing the relations in (3.18) and (3.19), we deduce that for  $x \geq 4$ ,

$$(3.20) \quad t_1 = x^{12}z^{36}\alpha(y)^3\alpha(w) < x^{12}z^{36}\eta_1\eta_2,$$

$$(3.21) \quad t_2 = (x^{10} - x^9 - 1)y^{36}w^{12}\beta(z)^3 > (x^{10} - x^9 - 1)y^{36}w^{12}\eta_3,$$

$$(3.22) \quad t_3 = x^7y^{12}z^{12}w^{12}\alpha(y)^2\beta(z)^2 > x^7y^{12}z^{12}w^{12}\eta_4\eta_5.$$

We continue to estimate  $e^{w+y-2z}$  and  $e^{z+x-2y}$ . Applying (3.9), (3.10), and (3.11) to  $w + y - 2z$ , we see that for  $x \geq 4$ ,

$$(3.23) \quad w + y - 2z < w_2 + y_2 - 2z_1,$$

which implies that

$$(3.24) \quad e^{w+y-2z} < e^{w_2+y_2-2z_1}.$$

In order to give a feasible upper bound for  $e^{w+y-2z}$ , we define

$$(3.25) \quad \Omega(t) = 1 + t + \frac{t^2}{2} + \frac{t^3}{6} + \frac{t^4}{24} + \frac{t^5}{120} + \frac{t^6}{720},$$

and it can be proved that for  $t < 0$ ,

$$(3.26) \quad e^t < \Omega(t).$$

Note that

$$w_2 + y_2 - 2z_1 = -\frac{\pi^4(108x^8 - 216\pi^2x^6 + 375\pi^4x^4 - 630\pi^6x^2 - 224\pi^8)}{972x^{11}} < 0$$

holds for  $x \geq 5$ , since

$$108x^8 - 216\pi^2x^6 > 0$$

for  $x > \sqrt{2}\pi \approx 4.443$ , and

$$375\pi^4x^4 - 630\pi^6x^2 - 224\pi^8 > 0$$

for  $x > \frac{\pi}{5}\sqrt{\sqrt{2443/3} + 21} \approx 4.422$ . Thus, by (3.26), we obtain that for  $x \geq 5$ ,

$$(3.27) \quad e^{w_2+y_2-2z_1} < \Omega(w_2 + y_2 - 2z_1).$$

Combining (3.24) and (3.27) yields that for  $x \geq 5$ ,

$$(3.28) \quad e^{w+y-2z} < \Omega(w_2 + y_2 - 2z_1).$$

Similarly, applying (3.9), (3.10), and (3.11) to  $z + x - 2y$ , we find that for  $x \geq 4$ ,

$$(3.29) \quad z_1 + x - 2y_2 < z + x - 2y,$$

so that

$$(3.30) \quad e^{z+x-2y} > e^{z_1+x-2y_2}.$$

Define

$$(3.31) \quad \omega(t) = 1 + t + \frac{t^2}{2} + \frac{t^3}{6} + \frac{t^4}{24} + \frac{t^5}{120} + \frac{t^6}{720} + \frac{t^7}{5040}.$$

It is true that for  $t < 0$ ,

$$(3.32) \quad \omega(t) < e^t.$$

We now give a lower bound for  $e^{z_1+x-2y_2}$ . Since

$$\begin{aligned} z + x - 2y &= \sqrt{x^2 + \frac{4\pi^2}{3}} + x - 2\sqrt{x^2 + \frac{2\pi^2}{3}} \\ &= \frac{-\left(\sqrt{x^2 + 4\pi^2/3} - x\right)^2}{\sqrt{x^2 + 4\pi^2/3} + x + 2\sqrt{x^2 + 2\pi^2/3}}, \end{aligned}$$

which is negative for  $n \geq 2$ , by (3.29), we deduce that for  $x \geq 4$ ,

$$(3.33) \quad z_1 + x - 2y_2 < 0.$$

Thus, applying (3.32) to (3.33) gives that for  $x \geq 4$ ,

$$(3.34) \quad e^{z_1+x-2y_2} > \omega(z_1 + x - 2y_2).$$

Combining (3.30) and (3.34), we find that for  $x \geq 4$ ,

$$(3.35) \quad e^{z+x-2y} > \omega(z_1 + x - 2y_2).$$

Using the above bounds for  $t_1, t_2, t_3, e^{w+z-2y}$  and  $e^{z+x-2y}$ , we obtain that for  $x \geq 5$ ,

$$(3.36) \quad \begin{aligned} &-e^{w+y-2z}t_1 + e^{z+x-2y}t_2 + 110t_3 \\ &> -\Omega(w_2 + y_2 - 2z_1)x^{12}z^{36}\eta_1\eta_2 \\ &\quad + \omega(z_1 + x - 2y_2)(x^{10} - x^9 - 1)y^{36}w^{12}\eta_3 + 110x^7y^{12}z^{12}w^{12}\eta_4\eta_5. \end{aligned}$$

To verify (3.7), we show that for  $x \geq 358$ ,

$$(3.37) \quad \begin{aligned} &-\Omega(w_2 + y_2 - 2z_1)x^{12}z^{36}\eta_1\eta_2 + \omega(z_1 + x - 2y_2)(x^{10} - x^9 - 1)y^{36}w^{12}\eta_3 \\ &\quad + 110x^7y^{12}z^{12}w^{12}\eta_4\eta_5 > 0. \end{aligned}$$

Substituting  $y, z$ , and  $w$  with  $\sqrt{x^2 + 2\pi^2/3}, \sqrt{x^2 + 4\pi^2/3}$ , and  $\sqrt{x^2 + 2\pi^2}$ , respectively, the left-hand side of the inequality (3.37) can be expressed as  $H(x)/G(x)$ , where

$$H(x) = \sum_{k=0}^{171} a_k x^k$$

and

$$G(x) = 39686201656473354776757087428535162639482880x^{88}.$$

Here we just list the values of  $a_{169}, a_{170}$ , and  $a_{171}$ :

$$\begin{aligned}
 a_{169} &= 734929660305062125495501619046947456286720 \\
 &\quad \times (35640 + 261360\pi^2 - 194\pi^6 - 249\pi^8), \\
 a_{170} &= 5879437282440497003964012952375579650293760 (7\pi^6 - 2970), \\
 a_{171} &= 4409577961830372752973009714281684737720320 (990 - \pi^6),
 \end{aligned}$$

which are all positive.

Given that  $G(x)$  is always positive, we aim to prove that  $H(x) > 0$ . Apparently,  $x \geq 2$  for  $n \geq 2$  and hence

$$(3.38) \quad H(x) \geq \sum_{k=0}^{170} -|a_k|x^k + a_{171}x^{171}.$$

Moreover, numerical evidence indicates that for any  $0 \leq k \leq 168$ ,

$$(3.39) \quad -|a_k|x^k > -a_{169}x^{169}$$

holds for  $x \geq 181$ . It follows that for  $x \geq 181$ ,

$$(3.40) \quad \sum_{k=0}^{170} -|a_k|x^k + a_{171}x^{171} > (-170a_{169} - a_{170}x + a_{171}x^2)x^{169}.$$

Combining (3.38) and (3.40), we obtain that for  $x \geq 181$ ,

$$(3.41) \quad H(x) > (-170a_{169} - a_{170}x + a_{171}x^2)x^{169}.$$

Thus,  $H(x)$  is positive provided

$$(3.42) \quad -170a_{169} - a_{170}x + a_{171}x^2 > 0,$$

which is true if

$$x > \frac{\sqrt{a_{170}^2 + 680a_{169}a_{171}} + a_{170}}{2a_{171}} \approx 357.867.$$

Hence we conclude that  $H(x)$  is positive when  $x \geq 358$ . This proves (3.37).

Combining (3.36) and (3.37), we find that for  $x \geq 358$ , or equivalently, for  $n \geq 19480$ , (3.7) holds, that is,

$$(3.43) \quad -e^{w+y-2z}t_1 + e^{z+x-2y}t_2 + 110t_3 > 0.$$

For  $2 \leq n \leq 19480$ , (3.43) can be directly verified. This completes the proof.  $\square$

#### 4. AN INEQUALITY ON $u_n$ AND $f(n)$

In this section, we present an inequality on  $u_n$  and  $f(n)$  that is also needed in the proof of Theorem 1.3.

**Theorem 4.1.** *Let  $u_n$  be defined as (1.19), that is,*

$$u_n = \frac{p(n+1)p(n-1)}{p(n)^2}.$$

For  $0 < t < 1$ , let

$$(4.1) \quad Q(t) = \frac{3t + 2\sqrt{(1-t)^3} - 2}{t^2}.$$

Then for  $n \geq 85$ ,

$$(4.2) \quad f(n) + \frac{110}{\mu(n-1)^5} < Q(u_n).$$

The proof of this theorem is based on Lemma 4.2, which gives an upper bound of  $f(n)$ . Recall that

$$f(n) = e^{x-2y+z} \frac{(x^{10} - x^9 - 1) y^{24} (z^{10} - z^9 - 1)}{x^{12} (y^{10} - y^9 + 1)^2 z^{12}},$$

where  $x, y, z, w$  are defined in (2.2).

**Lemma 4.2.** *Let  $\Omega(t)$  be defined as in (3.25), that is,*

$$\Omega(t) = 1 + t + \frac{t^2}{2} + \frac{t^3}{6} + \frac{t^4}{24} + \frac{t^5}{120} + \frac{t^6}{720},$$

and let  $y_1, y_2, z_1$ , and  $z_2$  be defined as in (3.12), (3.13), (3.14), and (3.15). For  $n \geq 4$ , we have

$$(4.3) \quad f(n) < \frac{\Omega(x - 2y_1 + z_2) y^{24} (x^{10} - x^9 - 1) (z^{10} - z^8 z_1 - 1)}{x^{12} z^{12} (y^{20} - 2y^{18} y_2 + y^{18} + 2y^{10} - 2y^8 y_2 + 1)} < 1.$$

*Proof.* To prove (4.3), we proceed to give estimates of the factors  $(y^{10} - y^9 + 1)^2$ ,  $z^{10} - z^9 - 1$ , and  $e^{x-2y+z}$  that appear in  $f(n)$ . The third inequality in (3.19) gives an estimate of  $(y^{10} - y^9 + 1)^2$ , that is, for  $x \geq 4$ ,

$$(4.4) \quad (y^{10} - y^9 + 1)^2 > y^{20} - 2y_2 y^{18} + y^{18} + 2y^{10} - 2y_2 y^8 + 1.$$

Using the bounds for  $y$  and  $z$  as given in (3.9) and (3.10), we are led to the following estimates for  $z^{10} - z^9 - 1$  and  $e^{x-2y+z}$  when  $x \geq 4$ ,

$$(4.5) \quad z^{10} - z^9 - 1 < z^{10} - z^8 z_1 - 1,$$

$$(4.6) \quad e^{x-2y+z} < e^{x-2y_1+z_2}.$$

To give an upper bound for  $e^{x-2y_1+z_2}$ , write

$$(4.7) \quad x - 2y_1 + z_2 = -\frac{\pi^4 (216x^8 - 216\pi^2 x^6 + 210\pi^4 x^4 - 210\pi^6 x^2 - 7\pi^8)}{1944x^{11}}.$$

For  $x > \pi$ , we have

$$216x^8 - 216\pi^2 x^6 > 0,$$

and for  $x > \pi \sqrt{(\sqrt{17/15} + 1)/2} \approx 3.192$ , we have

$$210\pi^4 x^4 - 210\pi^6 x^2 - 7\pi^8 > 0.$$

Therefore, it follows from (4.7) that for  $x \geq 4$ ,

$$x - 2y_1 + z_2 < 0,$$

which, together with (3.26), yields that for  $x \geq 4$ ,

$$(4.8) \quad e^{x-2y_1+z_2} < \Omega(x - 2y_1 + z_2).$$

Combining (4.6) and (4.8), we find that for  $x \geq 4$ ,

$$(4.9) \quad e^{x-2y+z} < \Omega(x - 2y_1 + z_2).$$

By means of the estimates in (4.4), (4.5), and (4.9), we arrive at the first inequality in (4.3).

To prove the second inequality in (4.3), recall the expressions of  $y$  and  $z$  in (3.8), namely,

$$y = \sqrt{x^2 + \frac{2\pi^2}{3}}, \quad z = \sqrt{x^2 + \frac{4\pi^2}{3}}.$$

It can be checked that

$$\Omega(x - 2y_1 + z_2)y^{24}(x^{10} - x^9 - 1)(z^{10} - z^8z_1 - 1) = \frac{I(x)}{x^{77}},$$

and

$$x^{12}z^{12}(y^{20} - 2y^{18}y_2 + y^{18} + 2y^{10} - 2y^8y_2 + 1) = N(x),$$

where  $I(x)$  is a polynomial in  $x$  of degree 121 and  $N(x)$  is a polynomial in  $x$  of degree 44. Thus we may assume that

$$\frac{\Omega(x - 2y_1 + z_2)y^{24}(x^{10} - x^9 - 1)(z^{10} - z^8z_1 - 1)}{x^{12}z^{12}(y^{20} - 2y^{18}y_2 + y^{18} + 2y^{10} - 2y^8y_2 + 1)} = \frac{K(x)}{J(x)},$$

where  $K(x)$  and  $J(x)$  are both polynomials of degree 121. Write

$$(4.10) \quad K(x) = \sum_{k=0}^{121} b_k x^k, \quad J(x) = \sum_{k=0}^{121} c_k x^k.$$

Here are the values of  $b_k$  and  $c_k$  for  $116 \leq k \leq 121$ :

$$b_{116} = -1398983398232765780459520\pi^4 (5181 + 41\pi^2),$$

$$b_{117} = 25181701168189784048271360\pi^2 (21 + 151\pi^2),$$

$$b_{118} = -4196950194698297341378560\pi^2 (258 + \pi^2),$$

$$c_{116} = -7197769583907579940464230400\pi^4,$$

$$c_{117} = 75545103504569352144814080\pi^2 (7 + 50\pi^2),$$

$$c_{118} = -1082813150232160714075668480\pi^2,$$

$$b_{119} = c_{119} = 12590850584094892024135680 (3 + 44\pi^2),$$

$$b_{120} = c_{120} = -75545103504569352144814080,$$

$$b_{121} = c_{121} = 37772551752284676072407040.$$

We claim that for  $x \geq 135$ ,

$$(4.11) \quad J(x) > 0,$$

and

$$(4.12) \quad J(x) - K(x) > 0.$$

It can be shown that for  $0 \leq k \leq 118$ ,

$$(4.13) \quad -|c_k|x^k > -c_{119}x^{119},$$



when

$$x > \pi \sqrt{\frac{6(7 + 50\pi^2)}{3 + 44\pi^2}} \approx 8.232.$$

It follows that for  $x \geq 9$ ,

$$(4.14) \quad J(x) > (-120c_{119} + c_{120}x + c_{121}x^2)x^{119}.$$

Since

$$(4.15) \quad -120c_{119} + c_{120}x + c_{121}x^2 > 0$$

when

$$x > 1 + \sqrt{11(11 + 160\pi^2)} \approx 133.255,$$

we find that  $J(x) > 0$  for  $x \geq 134$ .

Similarly, to prove (4.12), we observe that for  $0 \leq k \leq 115$ ,

$$(4.16) \quad -|c_k - b_k|x^k > -(c_{116} - b_{116})x^{116}$$

when

$$x > \frac{1}{2}\pi \sqrt{\frac{5616 + 3127\pi^2}{108 + 123\pi^2}} \approx 8.232.$$

Let

$$\theta(x) = -117(c_{116} - b_{116}) + (c_{117} - b_{117})x + (c_{118} - b_{118})x^2.$$

Then (4.16) implies that for  $x \geq 9$ ,

$$(4.17) \quad J(x) - K(x) = \sum_{k=0}^{118} (c_k - b_k)x^{118} > \theta(x)x^{116}.$$

Given that  $\theta(x)$  is positive when  $x \geq 135$ , we arrive at (4.12).

Combining (4.11) and (4.12), we deduce that the second inequality (4.3) is valid for  $x \geq 135$ , or equivalently, for  $n \geq 2771$ . The case for  $4 \leq n \leq 2771$  can be directly verified, and hence the proof is complete.  $\square$

We are now ready to prove Theorem 4.1.

*Proof of Theorem 4.1.* Recall that the theorem states that for  $n \geq 85$ ,

$$(4.18) \quad f(n) + \frac{110}{\mu(n-1)^5} < Q(u_n).$$

It can be checked that (4.18) is true for  $85 \leq n \leq 35456$ . We now show that (4.18) is true for  $n \geq 35457$ . Recall that

$$Q(t) = \frac{3t + 2\sqrt{(1-t)^3} - 2}{t^2},$$

and so

$$(4.19) \quad Q'(t) = \frac{t(\sqrt{1-t} - 3) - 4\sqrt{1-t} + 4}{t^3}.$$

Setting  $t = 1 - \tau$ , we get

$$Q'(t) = \frac{1}{(\sqrt{\tau} + 1)^3},$$

thus (4.19) can be rewritten as

$$(4.20) \quad Q'(t) = \frac{1}{(\sqrt{1-t} + 1)^3}.$$

As  $Q'(t)$  is positive for  $0 < t < 1$ ,  $Q(t)$  is increasing for  $0 < t < 1$ . By Theorem 2.1, we know that  $f(n) < u_n$  for  $n \geq 1207$ , so that for  $n \geq 1207$ ,

$$(4.21) \quad Q(f(n)) < Q(u_n).$$

Thus (4.18) is justified if we can prove that for  $n \geq 35457$ ,

$$(4.22) \quad f(n) + \frac{110}{\mu(n-1)^5} < Q(f(n)).$$

Let

$$(4.23) \quad \vartheta(t) = Q(t) - t = \frac{3t + 2\sqrt{(1-t)^3} - t^3 - 2}{t^2}.$$

In this notation, (4.22) says that for  $n \geq 35457$ ,

$$(4.24) \quad \vartheta(f(n)) > \frac{110}{\mu(n-1)^5}.$$

To prove the above inequality, we shall use the polynomials  $J(x)$  and  $K(x)$  as given by (4.10), namely,

$$K(x) = \sum_{k=0}^{121} b_k x^k, \quad J(x) = \sum_{k=0}^{121} c_k x^k.$$

Note that

$$(4.25) \quad \vartheta'(t) = -\frac{t^3 + 4(1-t)^{3/2} + 3t + 3t\sqrt{1-t} - 4}{t^3}.$$

Setting  $t = 1 - \tau$ , (4.25) becomes

$$\vartheta'(t) = -\frac{\sqrt{\tau}(\tau + 3\sqrt{\tau} + 3)}{(\sqrt{\tau} + 1)^3},$$

leading to the expression

$$\vartheta'(t) = -\frac{\sqrt{1-t}(3\sqrt{1-t} + 4 - t)}{(\sqrt{1-t} + 1)^3},$$

which is negative for  $0 < t < 1$ . Thus,  $\vartheta(t)$  is decreasing for  $0 < t < 1$ .

It can be seen from Lemma 4.2 that  $0 < f(n) < K(x)/J(x) < 1$  for  $n \geq 4$ , so that for  $n \geq 35457$ ,

$$(4.26) \quad \vartheta(f(n)) > \vartheta\left(\frac{K(x)}{J(x)}\right).$$

Because of (4.26), to verify (4.24), it is sufficient to show that for  $n \geq 35457$ ,

$$(4.27) \quad \vartheta\left(\frac{K(x)}{J(x)}\right) > \frac{110}{x^5}.$$

This goal can be achieved by finding an estimate for  $\vartheta(K(x)/J(x))$ . We first derive the following range of  $K(x)/J(x)$  for  $x \geq 134$ ,

$$(4.28) \quad \frac{\sqrt{5}-1}{2} < \frac{K(x)}{J(x)} < 1.$$

By Lemma 4.2, we know that  $K(x)/J(x) < 1$  for  $x \geq 4$  and  $J(x) > 0$  for  $x \geq 134$ . To justify (4.28), we only need to show that for  $x \geq 134$ ,

$$(4.29) \quad 2K(x) - (\sqrt{5}-1)J(x) > 0.$$

Note that

$$b_{119} = c_{119}, \quad b_{120} = c_{120}, \quad b_{121} = c_{121},$$

and it can be shown that for  $0 \leq k \leq 118$ ,

$$(4.30) \quad -|2b_k - (\sqrt{5} - 1)c_k|x^k > -(3 - \sqrt{5})c_{119}x^{119}$$

when

$$x > \pi \sqrt{\frac{\pi^2 (\sqrt{5} + 303) + 42}{3 + 44\pi^2}} \approx 8.303.$$

It follows that for  $x \geq 9$ ,

$$(4.31) \quad 2K(x) - (\sqrt{5} - 1)J(x) > (3 - \sqrt{5})(-120c_{119} + c_{120}x + c_{121}x^2)x^{119}.$$

Since

$$-120c_{119} + c_{120}x + c_{121}x^2 > 0$$

for  $x > \sqrt{11(11 + 160\pi^2)} + 1 \approx 133.255$ , we arrive at (4.29), and so (4.28) is proved.

The above range of  $K(x)/J(x)$  enables us to bound  $\vartheta(K(x)/J(x))$ . Recalling that

$$\vartheta(t) = \frac{3t + 2\sqrt{(1-t)^3 - t^3 - 2}}{t^2},$$

we obtain that

$$(4.32) \quad \vartheta(t) - (1-t)^{\frac{3}{2}} = -\frac{t^3 + t^2(1-t)^{\frac{3}{2}} - 2(1-t)^{\frac{3}{2}} - 3t + 2}{t^2}.$$

Set  $t = 1 - \tau$  to get

$$\vartheta(t) - (1-t)^{\frac{3}{2}} = -\frac{\tau^{\frac{3}{2}}(\tau + \sqrt{\tau} - 1)}{(\sqrt{\tau} + 1)^2}.$$

Thus

$$\begin{aligned} \vartheta(t) - (1-t)^{\frac{3}{2}} &= \frac{(1-t)^{\frac{3}{2}}(t - \sqrt{1-t})}{(\sqrt{1-t} + 1)^2} \\ &= \frac{(1-t)^{\frac{3}{2}}\left(t + \frac{\sqrt{5}+1}{2}\right)\left(t - \frac{\sqrt{5}-1}{2}\right)}{(\sqrt{1-t} + 1)^2(\sqrt{1-t} + t)}, \end{aligned}$$

which is positive for  $\frac{\sqrt{5}-1}{2} < t < 1$ , and hence, for  $\frac{\sqrt{5}-1}{2} < t < 1$  we have

$$(4.33) \quad \vartheta(t) > (1-t)^{\frac{3}{2}}.$$

In view of (4.28) and (4.33), we infer that for  $x \geq 134$ ,

$$(4.34) \quad \vartheta\left(\frac{K(x)}{J(x)}\right) > \left(1 - \frac{K(x)}{J(x)}\right)^{\frac{3}{2}}.$$

We continue to show that for  $x \geq 483$ ,

$$(4.35) \quad \left(1 - \frac{K(x)}{J(x)}\right)^{\frac{3}{2}} > \frac{110}{x^5}.$$

Since  $J(x) > 0$  for  $x \geq 134$ , the above inequality can be reformulated as follows. For  $x \geq 483$ ,

$$(4.36) \quad x^{10}(J(x) - K(x))^3 - 110^2 J(x)^3 > 0.$$

The left-hand side of (4.36) is a polynomial of degree 364, so that we may write

$$(4.37) \quad x^{10}(J(x) - K(x))^3 - 110^2 J(x)^3 = \sum_{k=0}^{364} \gamma_k x^k.$$

The values of  $\gamma_{364}$ ,  $\gamma_{363}$ , and  $\gamma_{362}$  are given below:

$$\gamma_{364} = 2^{72} 3^{105} 5^3 \pi^{12},$$

$$\gamma_{363} = -2^{73} 3^{107} 5^3 (490050 + \pi^{12}),$$

$$\gamma_{362} = 2^{72} 3^{105} 5^3 (52925400 + 144\pi^{12} + 41\pi^{14}).$$

For  $0 \leq k \leq 361$ , we find that

$$(4.38) \quad -|\gamma_k| x^k > -\gamma_{362} x^{362},$$

provided that

$$x > \frac{793881000 + 2328717600\pi^2 + 3996\pi^{12} + 4392\pi^{14} + \pi^{16}}{317552400 + 864\pi^{12} + 246\pi^{14}} \approx 20.126.$$

Thus, for  $x \geq 21$ ,

$$x^{10}(J(x) - K(x))^3 - 110^2 J(x)^3 > (-363\gamma_{362} + \gamma_{363}x + \gamma_{364}x^2)x^{362},$$

which is positive, since

$$-363\gamma_{362} + \gamma_{363}x + \gamma_{364}x^2 > 0$$

as long as

$$x > \frac{\sqrt{1452\gamma_{362}\gamma_{364} + \gamma_{363}^2} - \gamma_{363}}{2\gamma_{364}} \approx 482.959.$$

Hence (4.35) is confirmed. Combining (4.34) and (4.35), we are led to (4.27). The proof is therefore complete.  $\square$

### 5. PROOF OF THEOREM 1.3

In this section, we present a proof of Theorem 1.3 based on the intermediate inequalities in the previous sections. The theorem states that for  $n \geq 95$ ,

$$(5.1) \quad 4(1 - u_n)(1 - u_{n+1}) - (1 - u_n u_{n+1})^2 > 0,$$

where

$$u_n = \frac{p(n+1)p(n-1)}{p(n)^2}.$$

*Proof of Theorem 1.3.* We shall make use of the fact that  $u_n < 1$  for  $n \geq 26$ , as proved by DeSalvo and Pak [13]. In order to prove (5.1), we define

$$(5.2) \quad \varphi(t) = 4(1 - u_n)(1 - t) - (1 - u_n t)^2.$$

Then (5.1) says that for  $n \geq 95$ ,

$$(5.3) \quad \varphi(u_{n+1}) > 0.$$

For  $95 \leq n \leq 1206$ , (5.3) can be directly checked. We proceed to prove that (5.3) holds for  $n \geq 1207$ . Let  $Q(t)$  be as defined in (4.1), that is,

$$Q(t) = \frac{3t + 2\sqrt{(1-t)^3} - 2}{t^2}.$$

We claim that  $\varphi(t) > 0$  for  $u_n < t < Q(u_n)$ . Rewrite  $\varphi(t)$  as

$$\varphi(t) = -u_n^2 t^2 + (6u_n - 4)t - 4u_n + 3.$$

The equation  $\varphi(t) = 0$  has two real roots,

$$P(u_n) = \frac{3u_n - 2\sqrt{(1-u_n)^3} - 2}{u_n^2}, \quad Q(u_n) = \frac{3u_n + 2\sqrt{(1-u_n)^3} - 2}{u_n^2},$$

so that  $\varphi(t) > 0$  for  $P(u_n) < t < Q(u_n)$ . Furthermore, we see that

$$\varphi(u_n) = (1 - u_n)^3(u_n + 3) > 0,$$

which implies  $P(u_n) < u_n < Q(u_n)$ . Therefore,  $\varphi(t) > 0$  for  $u_n < t < Q(u_n)$ , as claimed.

To verify (5.3), it remains to show that for  $n \geq 1207$ ,

$$(5.4) \quad u_n < u_{n+1} < Q(u_n).$$

Recall that  $u_n < u_{n+1}$  holds for  $n \geq 116$ , as proved by Chen, Wang, and Xie [7]. By Theorem 2.1, we know that  $u_{n+1} < g(n+1)$  for  $n \geq 1207$ . But Theorem 3.1 asserts that for  $n \geq 2$ ,

$$g(n+1) < f(n) + \frac{110}{\mu(n-1)^5}.$$

Furthermore, Theorem 4.1 states that for  $n \geq 2$ ,

$$f(n) + \frac{110}{\mu(n-1)^5} < Q(u_n).$$

Thus we conclude that  $u_{n+1} < Q(u_n)$  for  $n \geq 1207$ , as claimed.  $\square$

#### ACKNOWLEDGMENTS

We wish to thank Ken Ono and the referee for insightful comments and suggestions.

#### REFERENCES

- [1] M. Aissen, I. J. Schoenberg, and A. M. Whitney, *On the generating functions of totally positive sequences. I* (English, with Hebrew summary), *J. Analyse Math.* **2** (1952), 93–103, DOI 10.1007/BF02786970. MR0053174
- [2] R. P. Boas Jr., *Entire functions*, Academic Press Inc., New York, 1954. MR0068627
- [3] G. Boros and V. Moll, *Irresistible integrals: Symbolics, analysis and experiments in the evaluation of integrals*, Cambridge University Press, Cambridge, 2004. MR2070237
- [4] P. Brändén, *Iterated sequences and the geometry of zeros*, *J. Reine Angew. Math.* **658** (2011), 115–131, DOI 10.1515/CRELLE.2011.063. MR2831515
- [5] W. Y. C. Chen, *Recent developments on log-concavity and q-log-concavity of combinatorial polynomials*, FPSAC 2010 Conference Talk Slides, <http://www.billchen.org/talks/2010-FPSAC.pdf> (2010).
- [6] W. Y. C. Chen, *The spt-function of Andrews*, *Surveys in Combinatorics 2017*, London Math. Soc. Lecture Note Ser., vol. 440, Cambridge Univ. Press, Cambridge, 2017, pp. 141–203. MR3728107
- [7] W. Y. C. Chen, L. X. W. Wang, and G. Y. B. Xie, *Finite differences of the logarithm of the partition function*, *Math. Comp.* **85** (2016), no. 298, 825–847, DOI 10.1090/mcom/2999. MR3434883
- [8] T. Craven and G. Csordas, *Jensen polynomials and the Turán and Laguerre inequalities*, *Pacific J. Math.* **136** (1989), no. 2, 241–260. MR978613
- [9] T. Craven and G. Csordas, *Problems and theorems in the theory of multiplier sequences*, *Serdica Math. J.* **22** (1996), no. 4, 515–524. MR1483603

- [10] G. Csordas, *Complex zero decreasing sequences and the Riemann hypothesis. II*, Analysis and applications—ISAAC 2001 (Berlin), Int. Soc. Anal. Appl. Comput., vol. 10, Kluwer Acad. Publ., Dordrecht, 2003, pp. 121–134, DOI 10.1007/978-1-4757-3741-7\_9. MR2022744
- [11] G. Csordas, T. S. Norfolk, and R. S. Varga, *The Riemann hypothesis and the Turán inequalities*, Trans. Amer. Math. Soc. **296** (1986), no. 2, 521–541, DOI 10.2307/2000378. MR846596
- [12] G. Csordas and R. S. Varga, *Necessary and sufficient conditions and the Riemann hypothesis*, Adv. in Appl. Math. **11** (1990), no. 3, 328–357, DOI 10.1016/0196-8858(90)90013-O. MR1061423
- [13] S. DeSalvo and I. Pak, *Log-concavity of the partition function*, Ramanujan J. **38** (2015), no. 1, 61–73, DOI 10.1007/s11139-014-9599-y. MR3396486
- [14] D. K. Dimitrov, *Higher order Turán inequalities*, Proc. Amer. Math. Soc. **126** (1998), no. 7, 2033–2037, DOI 10.1090/S0002-9939-98-04438-4. MR1459117
- [15] D. K. Dimitrov and F. R. Lucas, *Higher order Turán inequalities for the Riemann  $\xi$ -function*, Proc. Amer. Math. Soc. **139** (2011), no. 3, 1013–1022, DOI 10.1090/S0002-9939-2010-10515-4. MR2745652
- [16] I. Dolgachev, *Lectures on invariant theory*, London Mathematical Society Lecture Note Series, vol. 296, Cambridge University Press, Cambridge, 2003. MR2004511
- [17] S. Fisk, *Questions about determinants and polynomials*, arXiv:0808.1850 (2008).
- [18] M. Griffin, K. Ono, L. Rolén, and D. Zagier, *Jensen polynomials for Riemann’s zeta function and suitable arithmetic sequences*, preprint (2017).
- [19] G. H. Hardy and S. Ramanujan, *Asymptotic formulae in combinatory analysis*, Proc. London Math. Soc. (2) **17** (1918), 75–115, DOI 10.1112/plms/s2-17.1.75. MR1575586
- [20] D. Hilbert, *Theory of algebraic invariants*, Cambridge University Press, Cambridge, 1993. Translated from the German and with a preface by Reinhard C. Laubenbacher; Edited and with an introduction by Bernd Sturmfels. MR1266168
- [21] R. S. Irving, *Integers, polynomials, and rings: A course in algebra*, Undergraduate Texts in Mathematics, Springer-Verlag, New York, 2004. MR2025456
- [22] J. L. W. V. Jensen, *Recherches sur la théorie des équations* (French), Acta Math. **36** (1913), no. 1, 181–195, DOI 10.1007/BF02422380. MR1555086
- [23] S. Karlin, *Total positivity. Vol. I*, Stanford University Press, Stanford, Calif, 1968. MR0230102
- [24] O. M. Katkova, *Multiple positivity and the Riemann zeta-function*, Comput. Methods Funct. Theory **7** (2007), no. 1, 13–31, DOI 10.1007/BF03321628. MR2321798
- [25] J. P. S. Kung and G.-C. Rota, *The invariant theory of binary forms*, Bull. Amer. Math. Soc. (N.S.) **10** (1984), no. 1, 27–85, DOI 10.1090/S0273-0979-1984-15188-7. MR722856
- [26] H. Larson and I. Wagner, *Hyperbolicity of the partition Jensen polynomials*, preprint (2017).
- [27] D. H. Lehmer, *On the series for the partition function*, Trans. Amer. Math. Soc. **43** (1938), no. 2, 271–295, DOI 10.2307/1990043. MR1501943
- [28] D. H. Lehmer, *On the remainders and convergence of the series for the partition function*, Trans. Amer. Math. Soc. **46** (1939), 362–373, DOI 10.2307/1989927. MR0000410
- [29] B. Ja. Levin, *Distribution of zeros of entire functions*, Revised edition, Translations of Mathematical Monographs, vol. 5, American Mathematical Society, Providence, R.I., 1980. Translated from the Russian by R. P. Boas, J. M. Danskin, F. M. Goodspeed, J. Korevaar, A. L. Shields and H. P. Thielman. MR589888
- [30] P. A. MacMahon, *Combinatory analysis*, Two volumes (bound as one), Chelsea Publishing Co., New York, 1960. MR0141605
- [31] J. Mařík, *On polynomials, all of whose zeros are real* (Czech, with Russian and German summaries), Časopis Pěst. Mat. **89** (1964), 5–9. MR0180548
- [32] P. R. W. McNamara and B. E. Sagan, *Infinite log-concavity: developments and conjectures*, Adv. in Appl. Math. **44** (2010), no. 1, 1–15, DOI 10.1016/j.aam.2009.03.001. MR2552652
- [33] G. V. Milovanović, D. S. Mitrinović, and Th. M. Rassias, *Topics in polynomials: extremal problems, inequalities, zeros*, World Scientific Publishing Co., Inc., River Edge, NJ, 1994. MR1298187
- [34] V. H. Moll, *Combinatorial sequences arising from a rational integral*, Online J. Anal. Comb. **2** (2007), Art. 4, 17. MR2289956
- [35] C. P. Niculescu, *A new look at Newton’s inequalities*, J. Inequal. Pure Appl. Math. **1** (2000), no. 2, Article 17, 14. MR1786404
- [36] K. Ono, Private communication (2017).

- [37] N. Obrechhoff, *Zeros of polynomials*, Publ. Bulg. Acad. Sci., Sofia, 1963 (in Bulgarian), English translation by I. Dimovski and P. Rusev, The Marin Drinov Acad. Publ. House, Sofia, 2003.
- [38] G. Pólya, *Über die algebraisch-funktionentheoretischen Untersuchungen von J. L. W. V. Jensen*, Kgl. Danske Vid. Sel. Math.-Fys. Medd. **7** (1927) 3–33.
- [39] G. Pólya and J. Schur, *Über zwei Arten von Faktorenfolgen in der Theorie der algebraischen Gleichungen* (German), J. Reine Angew. Math. **144** (1914), 89–113, DOI 10.1515/crll.1914.144.89. MR1580897
- [40] H. Rademacher, *A convergent series for the partition function  $p(n)$* , Proc. Natl. Acad. Sci. USA, **23** (1937), no. 2, 78–84.
- [41] S. Rosset, *Normalized symmetric functions, Newton's inequalities and a new set of stronger inequalities*, Amer. Math. Monthly **96** (1989), no. 9, 815–819, DOI 10.2307/2324844. MR1033349
- [42] Q. I. Rahman and G. Schmeisser, *Analytic theory of polynomials*, London Mathematical Society Monographs. New Series, vol. 26, The Clarendon Press, Oxford University Press, Oxford, 2002. MR1954841
- [43] B. Sturmfels, *Algorithms in invariant theory*, Texts and Monographs in Symbolic Computation, Springer-Verlag, Vienna, 1993. MR1255980
- [44] G. Szegő, *On an inequality of P. Turán concerning Legendre polynomials*, Bull. Amer. Math. Soc. **54** (1948), 401–405, DOI 10.1090/S0002-9904-1948-09017-6. MR0023954
- [45] G. Szegő, *Orthogonal Polynomials*, American Mathematical Society, New York, 1939. American Mathematical Society Colloquium Publications, v. 23. MR0000077
- [46] C. G. Wagner, *Newton's inequality and a test for imaginary roots*, Two-Year College Math. J. **8** (1977), no. 3, 145–147.

CENTER FOR COMBINATORICS, LPMC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA; AND CENTER FOR APPLIED MATHEMATICS, TIANJIN UNIVERSITY, TIANJIN 300072, PEOPLE'S REPUBLIC OF CHINA

*Email address:* chen@nankai.edu.cn

CENTER FOR COMBINATORICS, LPMC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

*Email address:* dennisjq@mail.nankai.edu.cn

CENTER FOR COMBINATORICS, LPMC, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

*Email address:* wsw82@nankai.edu.cn